

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
3 October 2002 (03.10.2002)

PCT

(10) International Publication Number
WO 02/077023 A2(51) International Patent Classification⁷: **C07K 14/435**(21) International Application Number: **PCT/CH02/00063**

(22) International Filing Date: 1 February 2002 (01.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/277,976 23 March 2001 (23.03.2001) US(71) Applicant (for all designated States except US): **UNIVERSITÄT ZÜRICH [CH/CH]**; Rämistrasse 71, CH-8006 Zürich (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KRAMPS, Thomas [DE/CH]**; Dorfstrasse 51, CH-8037 Zürich (CH). **BASLER, Konrad [CH/CH]**; Traubenweg 34, CH-8700 Küsnacht (CH).(74) Agent: **LIEBETANZ, Michael**; Isler & Pedrazzini AG, Postfach 6940, CH-8023 Zürich (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,

CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A NEW ESSENTIAL DOWNSTREAM COMPONENT OF THE WINGLESS SIGNALLING PATHWAY

(57) Abstract: The present invention relates to a new essential downstream component of the wingless signaling pathway. In particular, the invention relates to nucleotide sequences of the *Drosophila melanogaster* daughter of legless (*doll*) gene, of its encoded proteins, as well as derivatives, fragments and analogues thereof. The invention includes vertebrate and invertebrate homologues of the Doll protein, comprising proteins that contain a stretch of amino acids with similarity to the *Drosophila* Doll gene. Methods for producing the Doll protein, derivatives and analogs, e.g. by recombinant means, and antibodies to Doll are provided by the present invention as well. The invention also relates to methods for performing high throughput screening assays for compounds modulating Doll function in the Wnt pathway.

A NEW ESSENTIAL DOWNSTREAM COMPONENT OF THE WINGLESS SIGNALLING PATHWAY

The present invention relates to a new essential downstream component of the wingless signaling pathway. In particular, the invention relates to nucleotide sequences of the *Drosophila melanogaster* daughter of legless (*doll*) gene, of its encoded proteins, as well as derivatives, fragments and analogues thereof. The invention includes vertebrate and invertebrate homologues of the Doll protein, comprising proteins that contain a stretch of amino acids with similarity to the *Drosophila* Doll gene. Methods for producing the Doll protein, derivatives and analogs, e.g. by recombinant means, and antibodies to Doll are provided by the present invention as well. The invention also relates to methods for performing high throughput screening assays for compounds modulating Doll function in the Wnt pathway.

Background of the invention

Wnt genes encode a large family of secreted, cysteine rich proteins that play key roles as intercellular signaling molecules in a wide variety of biological processes (for an extensive review see (Wodarz and Nusse 1998). The first Wnt gene, mouse *wnt-1*, was discovered as a proto-oncogene activated by integration of mouse mammary tumor virus in mammary tumors (Nusse and Varmus 1982). Consequently, the involvement of the Wnt pathway in cancer has been largely studied. With the identification of the *Drosophila* polarity gene *wingless* (*wg*) as a *wnt-1* homologue (Cabrera, Alonso et al. 1987; Perrimon and Mahowald 1987; Rijsewijk, Schuermann et al. 1987), it became clear that *wnt* genes are important developmental regulators. Thus, although at first glance dissimilar, biological processes like embryogenesis and carcinogenesis both rely on cell communication via identical signaling pathways. In a current model of the pathway, the secreted Wnt protein binds to Frizzles cell surface receptors and activates the cytoplasmic protein Dishevelled (Dsh). Dsh then transmits the signal to a complex of several proteins, including

the protein kinase Shaggy/GSK3, the scaffold protein Axin and β -Catenin, the vertebrate homologue of Armadillo. In this complex β -Catenin is targeted for degradation after being phosphorylated by Sgg. After Wnt signaling and the resulting down-regulation of Sgg activity, β -Catenin (or its *Drosophila* homologue Armadillo) escape from degradation and accumulate into the cytoplasm. Free cytoplasmic β -Catenin translocates to the nucleus by a still obscure mechanism, and modulates gene transcription through binding the Tcf/Lef family of transcription factors (Grosschedl R 1999).

This set up, in which the key transducer is continuously held in check, is highly susceptible to mutations in its inhibitory components. The loss of any of the three elements of the β -Catenin destruction complex leads to an increase in β -Catenin levels, and hence to the constitutive activation of the pathway. While this may reduce cellular viability, as upon loss of GSK-3 function, it can also lead to cell fate changes, uncontrolled proliferation and tumorigenic behavior as in the cases of APC and Axin (Barker N 1999; Morin 1999; Potter 1999; Roose and Clevers 1999; Waltzer and Bienz 1999). Attempts to counter these harmful situations must aim at curbing the nuclear activities of β -Catenin, either by preventing the formation of the β -Catenin-TCF complex or by interfering with its transcriptional activator function.

Currently, there are no known therapeutic agents effectively inhibiting β -Catenin transcriptional activation. This is partly due to the fact that many of the essential components required for its full activation and nuclear translocation are still unknown. Consequently, there is an urge to understand more about this pathway in order to be able to develop effective drugs against these highly malignant diseases.

In order to identify new components required for Wingless activation the inventors used a *Drosophila* genetic approach to screen for dominant suppressors of the rough eye phenotype caused by ectopic expression of Wingless, the *Drosophila* homologue of Wnt, during eye development. Three genes were identified: the β -catenin homologue *armadillo* (*arm*), the *tcf/lef-1* homologue *pangolin* (*pan*) and *legless* (*lgs*), a completely new gene (US 09/915.543). The *lgs* gene was subsequently cloned and

its *in vivo* requirement for Wingless signal transduction in embryo and in developing tissues was confirmed. The presence of Lgs is required for a transcriptional active Arm/Pangolin complex and over-expression of Lgs strongly stimulates the transcriptional output of this bipartite transcription factor. The human genome contains at least two human Lgs homologues. One of them, Bcl9, has been previously implicated in B cell malignancies (Willis, Zalberg et al. 1998). It was also genetically and biochemically demonstrated that dLgs and hLgs bind to Armadillo and β -Catenin and are functionally required for Wnt signal propagation in human cells. However, genetic experiments strongly suggested the presence of a second protein which binds to Lgs and is essential for the function of the active β -Catenin-Pangolin-Lgs complex.

The present invention describes the cloning and functional characterization of a novel *Drosophila* protein, named Daughter of Legless (Doll), which binds to Lgs and is required for Wnt signaling. In addition, the invention provides the sequences of the functional and structural human and mouse homologues as well as methods to screen for compounds inhibiting Doll function in the Wnt pathway.

Definitions

The term "Doll polypeptide", "Doll protein" when used herein encompasses native invertebrate and vertebrate Doll and Doll variant sequences (which are further defined herein).

A "wild type sequence Doll" comprises a polypeptide having the same amino acid sequence as a Doll protein derived from nature. Such wild type sequence of Doll can be isolated from nature or produced by recombinant and/or synthetic means. The term "wild type sequence Doll" specifically encompasses naturally occurring truncated forms, naturally occurring variant forms (e.g., alternatively spliced forms) and naturally occurring allelic variants of Doll. In one embodiment of the invention, the wild type Doll sequence is a mature or full-length Doll sequence comprising amino acids 1 to 815 of dDoll (Figure 1), or 1 to 419 of hDoll-1, or 1 to 406 of hDoll-2 (Figure 2), or 1 to 417 of mDoll-1, or 1 to 407 of mDoll-2 (Figures 3).

"Doll variant" means an active Doll, having at least about 50% amino acid sequence identity with the amino acid sequence of a wild type Doll protein of Figure 1, 2 and 3. The term "Doll variant" however, does also include functional homologues of Doll in the Wnt pathway.

"Percent (%) amino acid sequence identity" with respect to the Doll sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the Doll sequence, after aligning the sequence and introducing gaps, if necessary, to achieve the maximum percentage sequence identity, and not considering any conservative amino acid substitution as part of the sequence identity. The % identity values used herein can be generated by WU-BLAST-2, which was obtained from (Tatusova TA 1999). WU-BLAST-2 uses several search parameters, most of which are set to the default values.

The term "positive", in the context of sequence comparison performed as described above, includes residues in the sequence compared that are not identical but have similar properties (e.g. as a result of a conservative substitution). The % value of positive is determined by the fraction of residues scoring a positive value in the BLOSUM 62 matrix divided by the total number of residues in the longer sequence as defined above.

In a similar manner, "percent (%) nucleic acid sequence identity" with respect to the coding sequence of the Doll polypeptides identified herein is defined as the percentage of nucleotide residues in a candidate sequence that are identical with the nucleotide residues in any of the Doll coding sequences of this invention. The identity values used herein can be generated using BLAST module of WU-BLAST-2 set to the default parameters.

The term "epitope tag" refers to a chimeric polypeptide comprising a Doll polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough that it does not interfere with the activity of the Doll polypeptide to which it is fused.

Nucleic acids are "operably linked" when are placed in a functional relationship with another nucleic acid sequence.

The term „epistasis“ means hierarchy in gene action. Epistasis experiments are performed to place components of a signaling pathway in the right order.

The term „rescue experiments“ are designed to determine which gene is responsible for a specific mutant phenotype. Specifically, mutant embryos are injected with coding or genomic DNA, and the effect of the introduced DNA is determined on the basis of the capacity to revert the mutant phenotype.

„Active“ or „activity“ refers to forms of Doll polypeptides that retain the biological and/or immunological activity. A preferred activity includes for instance the ability to modulate the Wnt signaling pathway.

The term „antagonist“ is used in a broad sense, and includes any molecule that partially or fully inhibits, blocks or neutralizes a biological activity of Doll polypeptides described herein. In a similar way, the term „agonist“ is used in the broadest sense and includes any molecule that mimics or support a biological activity of an active Doll polypeptide.

„Treatment“ refers to both therapeutic treatments and prophylactic or preventive measures, wherein the objective is to prevent or slow down the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

Summary of the invention

The present invention relates to a novel family of proteins present in insects and vertebrate organisms, referred to hereinafter as „Daughter of Legless (Doll)“ proteins. These proteins play an essential role in the Wnt signaling pathway, and thus in the formation and maintenance of spatial arrangements and proliferation of tissues during development, and in the formation and growth of many human tumors.

In particular, the invention relates to nucleotide sequences of the *Drosophila melanogaster* *doll* gene, of proteins encoded by said nucleotide sequences, as well as fragments, derivatives and structural and functional analogs thereof.

In a preferred embodiment the invention relates to the nucleotide and protein sequences of the human and mouse *doll* homologues, *hdoll-1*, *hdoll-2* and *mdoll-1* and *mdoll-2*, respectively.

In one embodiment, the isolated nucleic acid comprises a sequence encoding a polypeptide having at least 50% amino acid sequence identity, preferably at least about 70% sequence identity, more preferably at least 90% sequence identity, even more preferably at least about 95% sequence identity, yet even more preferably at least about 98% sequence identity, and most preferably 100% identity to (a) a fragment or the entire protein sequence of the Doll polypeptide shown in Figure 1, or (b) the complement of the nucleic acid molecule coding for (a).

In another preferred embodiment, the isolated nucleic acid encodes a polypeptide having at least 50% amino acid sequence identity, preferably about 70% sequence identity, more preferably at least 90% sequence identity, even more preferably about 95% sequence identity, yet even more preferably about 98% sequence identity, and most preferably 100% identity to (a) a polypeptide which is part or the entire human Doll polypeptides of figure 2a/b or (b) the complement of the nucleic acid molecule coding for (a).

In another embodiment, the isolated nucleic acid encodes a polypeptide sequence having at least 50% amino acid sequence identity, preferably about 70% sequence identity, more preferably at least 90% sequence identity, even more preferably about 95% sequence identity, yet even more preferably about 98% sequence identity, and most preferably 100% identity to (a) a polypeptide encoding part of the entire mouse Doll protein of figure 3 a/b or (b) the complement of the nucleic acid molecule coding for (a).

In a further embodiment, the isolated nucleic acid comprises a sequence encoding a polypeptide with a low overall amino acid sequence identity but shows a sequence identity of at least 60%,

preferably at least 70%, more preferably at least 80%, even more preferably at least 90% and most preferably 100% in the conserved domains DHD and PHD (Figure 4).

In yet another embodiment of the present invention isolated nucleic acids encode polypeptides having a function resembling that of the *doll* genes.

In another embodiment, the invention relates to a fragment of the *Drosophila* or vertebrate *doll* nucleic acid sequences that is applied as hybridization probe. Such nucleic acid fragments are about 20 to about 100 nucleotides in length, preferably from about 20 to about 60 nucleotides in length, most preferably from 20 to 50 nucleotides in length and are derived from the nucleotides sequences shown in Figure 1, 2 and 3.

The invention further provides eucaryotic and procaryotic expression vectors comprising a nucleic acid molecule encoding *Drosophila* or vertebrate *doll* or a fragment thereof as shown in Figures 1, 2 and 3. The vector can comprise any of the molecules or fragments thereof described above.

The invention also includes host cells comprising such a vector. By way of example, the host cells can be mammalian cells, yeast cells, insect cells, plant cells or bacteria cells.

Methods of production, isolation and purification of the Doll proteins, derivatives and analogs, e.g. by recombinant means, are also provided (see Example VI, below). In a specific aspect, the invention concerns an isolated Doll peptide, comprising an amino acid sequence of at least 80%, preferably at least about 85% sequence identity, more preferably at least 90% sequence identity, even more preferably at least 95% sequence identity, yet most preferably 100% identity with the amino acid sequences of Figures 1, 2 and 3.

In yet another embodiment the invention relates to chimeric proteins comprising a Doll polypeptide fused to a heterologous polypeptide or amino acid sequence. An example of such chimeric molecule comprises a Doll polypeptide fused to an epitope tagged sequence, glutathione-S-transferase protein or to a protein with

In another embodiment the invention provides antibodies that specifically recognize Doll polypeptides. The antibodies can be a polyclonal or a monoclonal preparation or fragments thereof. Polyclonal antibodies are prepared by immunization of rabbits with purified Doll polypeptides prepared as described in Example VI.

In yet another embodiment, the invention also features animals, e.g. *Drosophila*, mice, rats, chicken, frogs, pigs or sheep, having a mutation in the *doll* gene, e.g. deletions, point mutations, foreign DNA insertions or inversions. Such animals can serve to study diseases characterized by disrupted Wnt function or in drug screening.

RNSDOCID: 6WO 0207702342 1 >

thereto, *doll* antisense DNA or RNA, *doll* double stranded RNA, and any chemical or natural occurring compound interfering with *Doll* function, synthesis or degradation. In particular, the invention provides methods to screen for chemical compounds, organic products or peptides interfering with *Doll* function in the Wnt pathway. In a preferred embodiment the screening method will be a cellular reporter gene assay or a protein-protein interaction assay.

In another embodiment, a screening assay based on protein-protein interaction is used to screen for compounds specifically inhibiting *Doll*-Lgs or *Doll*-interaction partner X.

The invention also provides methods to screen for chemical compounds, organic products or peptides interfering with *Doll* function in the Wnt pathway.

Furthermore, the invention comprises the use of the DHD domain in screening assays such as an *in vitro* protein-protein interaction assay or a protein-protein interaction in a host cell. Said assays are applied for the identification of chemical compounds, organic products, polypeptides or peptides interfering with *Doll* function in the Wnt pathway.

In a preferred embodiment, a therapeutic product according to the invention is administered to treat a cancerous condition or to prevent progression from a pre-neoplastic or non-malignant condition to a neoplastic or malignant state.

In other specific embodiments, a therapeutic product of the invention is administered to treat a blood disease or to promote tissue regeneration and repair. Finally disorders of cell fate, especially hyperproliferative or hypoproliferative disorders, involving aberrant or undesirable expression, or localization, or activity of the *Doll* protein can be diagnosed by detecting such levels.

Brief description of the drawings

Figure 1 The *Drosophila* *doll* cDNA and protein sequence.

Figure 2 The human *doll-1* and *doll-2* cDNA and protein sequence.

Figure 3 The mouse *doll-1* and *doll-2* cDNA and protein sequences.

Figure 4. *Drosophila* and human Doll proteins contain a PHD finger motif with which they bind to the HD1 of Lgs/BCL9

(A) Top: Schematic representation of Doll, human DOLL-1 and human DOLL-2. The two domains that show high sequence similarities are highlighted in dark gray (DHD: Doll homology domain) and red (PHD: plant homology domain). The DHD appears to be unique, as the inventors failed to find a similar sequence in other *Drosophila* or human proteins. GenBank accession numbers for Doll, hDoll-1, hDoll-2 are AF457206, AF457207, AF457208, respectively. Bottom: Multiple alignment of *Drosophila*, human and mouse Doll protein sequences.

(B) Alignment of amino acid sequences of DHD and PHD in Doll and its human homologues.

Similarities are boxed, identities shaded in gray. The numbers to the left indicate the positions of DHD and PHD within their respective protein sequences. For the DHD alignment a gap of 22 aa has been introduced in the *Drosophila* DHD (represented as (X)5).

(C) Mapping of the Lgs/BCL9 interaction site in Doll. Schematic representation of the proteins tested in the yeast-two-hybrid assay for their interactions with Lgs and BCL9. Results are indicated to the right ("bdg").

(D) Mapping of the Doll interaction site in dLgs and hLgs/BCL9. Schematic representation of the proteins tested in the yeast-two-hybrid assay for their interactions with Lgs. The two proteins shown at the bottom were tested by a pull-down assay for both dLgs (numbers without brackets) and hLgs/BCL9 (numbers in parenthesis) with the same result ("bdg"). The deletion removing HD1 comprises aa 318-345 for Lgs and aa 177-204 for hLgs/BCL9. Fusion proteins used were S-Tag-dDoll (aa 542-815) and GST-hDOLL-2 (aa 301-406).

Figure 5. *doll* is a segment polarity gene required for Wg signalling

(A-C) Cuticle preparations of larvae derived from wild-type (A), *wg* mutant (B), and *doll* mutant embryos (C). The *doll*¹³⁰/*doll*¹³⁰ embryo in (C) is derived from a homozygous *doll*¹³⁰ mutant germ line clone (see Experimental Procedures) and displays a *wg*-like phenotype.

(D,E) *doll* functions downstream of dAPC2. Two cuticle preparations are shown from larvae that developed in the absence of maternal and zygotic wild-type dAPC2 function (McCartney et al., 1999). The embryo in (E) additionally lacks the maternal and zygotic function of *doll* (see Examples). In contrast to dAPC single mutant animals, which have strongly reduced denticle belts, double mutants display a *doll*-like phenotype.

(F-I) Confocal images of third instar wing imaginal disc preparations stained with antibodies against Doll (F,G) and Ptc (H,I). Wild-type animals show normal expression of these genes (F,H). Discs derived from *doll*¹³⁰ mutant larvae are small, yet express Ptc (I), but fail to express Dll (H). Lack of Dll expression may be an indirect consequence of the earlier wing-to-notum transformation in *doll*¹³⁰ larvae. However, we also see a strong reduction of Dll expression in *doll*¹³⁰ mutant cells from mosaic animals (not shown).

Figure 6. Lgs and the PHD finger of Doll serve to assemble Doll and Arm

Schematic representation of Lgs (yellow) and Doll (light green) constructs that were used in transgene assays to assess their ability to rescue *lgs* or *doll* mutant animals. 1: Full-length Lgs (pOP216, aa 1-1464). 2: C-terminally truncated Lgs (pTK131, aa 1-583). 3: HD1-Gal11-HD2 (pTK153, aa 268-395 (HD1), aa 369-500 (Gal11), aa 465-596 (HD2)). 4: HD1-(HA)3-HD2 (pTK143, aa 268-395 (HD1), aa 465-596 (HD2)). 5: Full-length Doll (pTK56, aa 1-815). 6: Doll[DPHD]-HD2 (pTK135, aa 1-740 (Doll[DPHD]), aa 483-561 (HD2)). Transgenes 1 to 4 are

able to rescue *lgs*^{20F} homozygotes. An example for an adult animal rescued by transgene 3 is shown on the right. Transgene 5 can rescue *doll*¹³⁰ homozygotes. Transgene 6 can rescue *doll*¹³⁰ as well

as *lgs*^{20F} homozygotes (photographs on the right).

Figure 7: Rescue of *ddoll*^{-/-} flies by expression of a human *doll* transgene. The lethality caused by the *doll*¹³⁰/EP(3)1076 genotype can be fully rescued by a tubulin 1 promoter-driven transgene that contains either the coding region of the *Drosophila* *doll* gene (not shown) or that of one of its two human homologues *hdoll*-1 and *hdoll*-2.

Figure 8: Effects of human *Doll* 1 and 2 on *Tcf* transcription. 293 cells were transiently transfected with the pTOPFLASH or pFOPFLASH luciferase reporters and different effector plasmids as indicated. A constitutively active form of β -Catenin (Δ N- β -Catenin, 50 ng) or human *Doll*-1 or *hDoll*-2 (350 ng) activate the pTOPFLASH reporter. Cotransfection of human *Doll* with Δ N- β -catenin strongly enhance the response.

Detailed description of the invention

The Wnt signaling cascade is essential for the development of both invertebrates and vertebrates, and has been implicated in tumorigenesis. The *Drosophila* *wg* genes are one of the best characterized within the Wnt-protein family, which includes more than hundred genes. In the *Drosophila* embryo, *wg* is required for formation of parasegment boundaries and for maintenance of *engrailed* (*en*) expression in adjacent cells. The epidermis of embryo defective in *wg* function shows only a rudimentary segmentation, which is reflected in an abnormal cuticle pattern. While the ventral cuticle of wild type larvae displays denticle belts alternating with naked regions, the cuticle of *wg* mutant larvae is completely covered with denticles. During imaginal disc development, *wg* controls dorso-ventral positional information. In the leg disc, *wg* patterns the future leg by the induction of ventral fate (Struhl and Basler 1993). In animals with reduced *wg* activity, the ventral half of the leg develops into a mirror image of the dorsal side (Baker 1988). Accordingly, reduced *wg* activity leads to the transformation of wing to notal tissue, hence the name of the gene (Sharma and Chopra 1976). In the eye disc, *wg* suppresses ommatidial differentiation in favor of head cuticle development, and is involved in establishing the dorso-

ventral axis across the eye field (Heberlein, Borod et al. 1998).

Additional genes have been implicated in the secretion, reception or interpretation of the Wg signaling. For instance, genetic studies in *Drosophila* revealed the involvement of *frizzled* (*Dfz*), *Dishevelled* (*dsh*), *shaggy/zeste-white-3* (*sgg/zw-3*), *armadillo* (*arm*), *adenomatous polyposis coli* (*E-apc*), *axin*, and *pan-golin* (*pan*) in Wg signaling. The genetic order of these transducers has been established in which Wg acts through Dsh to inhibit Sgg, thus relieving the repression of Arm by Sgg, resulting in the cytoplasmic accumulation of Arm and its translocation to the nucleus. In the nucleus Arm interacts with Pan to activate transcription of target genes. Vertebrate homologues have been identified for all these components (for an updated review see (Peifer and Polakis 2000), suggesting that novel identified members of the *Drosophila* signaling pathway may likely have vertebrate counterparts.

Mutations leading to nuclear accumulation of the mammalian homologue of Arm, β -Catenin, and consequently to constitutive activation of the Wnt pathway have been observed in many types of cancer, including colon cancer, breast cancer, melanoma, hepatocellular carcinoma, ovarian cancer, endometrial cancer, medulloblastoma pilomatricomas, and prostate cancer (Morin 1999; Polakis, Hart et al. 1999). It is now apparent that deregulation of β -Catenin signaling is an important event in the genesis of these malignancies. However, there are still no known therapeutic agents effectively inhibiting β -Catenin transcriptional activation. This is partly due to the fact that many of the essential components required for its full activation and nuclear translocation are still unknown.

In order to identify new components required for Wingless activation the inventors used a *Drosophila* genetic approach to screen for dominant suppressors of the rough eye phenotype caused by ectopic expression of Wingless (Wg), the *Drosophila* homologue of Wnt, during eye development. A new gene, *legless* (*lgs*, US09/915.543) was identified as a strong dominant suppressor of the rough eye phenotype. The gene was subsequently cloned and its *in vivo* requirement for Wg signal transduction in embryo and in developing tissues was confirmed. The human genome contains at least two human Lgs homologues, hLgs/Bcl9 and hLgs-1.

dLgs and hLgs bind to Armadillo and β -Catenin and are functionally required for Wnt signal propagation in invertebrate and vertebrate cells (US 09/915.543). In particular, the presence of Lgs is required for a transcriptional active Arm/Pangolin complex and over-expression of Lgs strongly stimulates the transcriptional output.

The inventors later made the interesting observation that a mutant form of Lgs protein from which the β -Catenin interacting domain was deleted exhibited a strong dominant-negative effect on Wg-dependent patterning processes when expressed from a transgene in wild-type larvae (data not shown). This strongly suggested that Lgs normally interacts not only with Arm, but also with at least one additional component. In an effort to identify such components yeast-two-hybrid screens for interacting proteins were carried out. In two independent screens in which either the entire protein or the N-terminal half of Lgs was used as a bait, a novel PHD finger protein, referred to as Daughter-of-Legless (Doll), was identified as a Lgs binding protein (Figure 4). The 815 amino acid residue Doll protein carries a C-terminal domain of 60 amino acids (Figure 4a), which shows extensive homologies to the PHD (plant homology domain) finger, also known as LAP (leukemia associated protein) domain (Aasland, Gibson et al. 1995). This domain comprises a cysteine rich Zn-binding motif, that has been associated with proteins involved in chromatin-mediated regulation of transcription. The PHD finger of Doll is necessary and sufficient to mediate the interaction to Lgs (Figure 4c-d). The inventors also demonstrate herein that this interaction is essential for Doll function.

The region of Lgs responsible for Doll-binding was mapped to the HD1 sequence (Figure 4d). Moreover, two human homologues of the *Drosophila* doll gene were identified and isolated (Figure 4a). The protein products of both human genes, hDOLL-1 and hDOLL-2, as well as their mouse homologues possess a highly conserved PHD finger which interacts with the HD1 of hLgs/BCL9 (Figure 4d). The only other domain in *Drosophila* Doll, hDoll-1/hDoll-2 and mDoll-1/mDoll-2 that shows significant sequence homology is a 50 amino acid stretch in the N-terminal region, which is referred herein to as 'Doll homology domain' (DHD, Figure 4a,b).

The interaction with Doll appears to be relevant for the *in vivo* function of Lgs, since a mutant form of Lgs with a deletion of

HD1 was unable to rescue *lgs* mutant animals. The physical association of *Doll* and *Lgs* suggested that *Doll*, like *Lgs*, may be required for *Wg* signaling in vivo. To explore this hypothesis a proprietary collection of suppressors of the *sev-wg* phenotype was searched for mutations that map to the tip of the right arm of chromosome 3, the position of the *doll* gene. One such suppressor, *Sup*¹³⁰, mapped to this position, and intriguingly, it showed dominant lethality in combination with the *lgs* allele *lgs*^{17E} (US09/915.543) (*Sup*¹³⁰ /+ *lgs*^{17E} /+ transheterozygous animals do not survive). The *doll* coding region was sequenced using genomic DNA from homozygous *Sup*¹³⁰ mutant larvae and a 14 bp deletion starting at amino acid position 751 was identified. Hence this allele is referred to as *doll*¹³⁰ and encodes a truncated *Doll* protein lacking the C-terminal PHD finger.

The lethality caused by the homozygous *doll*¹³⁰ genotype can be fully rescued by a tubulin 1 promoter-driven transgene that contains either the coding region of the *Drosophila* *doll* gene or that of one of its two human homologues *hDoll-1* and *hDoll-2* (Figure 7). Thus, the vertebrate homologues of *doll* were confirmed genetically to be true functional homologues of *Doll*, and hence the vertebrate homologues are part of this invention.

To assay the possible role of *Doll* in *Wg* signal transduction during development, embryos homozygous for the *doll*¹³⁰ mutation that derived from female germ cells equally mutant for *doll* were generated. *Doll* mRNA is maternally contributed and strongly and ubiquitously expressed during all the developmental stages. Consequently, only embryos lacking both embryonal and maternal *doll* are characterized by a severe segment polarity phenotype (Figure 5A-C), while weaker loss of function *doll* mutants display pupal lethality with a partial or complete loss of the antennae and the legs. Mutant individuals lacking only zygotic function survive until early pupal stages and exhibit imaginal discs that are abnormally small. The Hh target gene *ptc* was expressed at wild-type levels in these discs, however, no expression of the *Wg* target *Dll* could be detected (Figure 5F-I). These discs appear to lack the presumptive wing blade field and possess two primordia for the notum (Figures 5G and I). The fact that similar phenotypes are caused by loss of function of *wg*, *dsh*, *arm*,

or *lgs* confirms the essential role of *doll* in the Wg signaling pathway.

To address the role of Doll in β -Catenin-mediated transcription a TCF reporter gene (TOPFLASH, (Morin, Sparks et al. 1997)) was used in immortalized human embryo kidney cells (HEK 293 cells). Low levels of a stable mutant form of β -Catenin (ΔN - β -catenin; (van de Wetering, Cavallo et al. 1997)) were introduced into these cells to partially stimulate the pathway. The additional expression of hDoll-1 (Figure 8) or hDoll-2 (not shown) lead to a large increase in luciferase activity (30-fold). These levels are significantly higher than the sum of those produced by either treatment alone (Figure 8). This potentiation of β -Catenin activity by hDoll-1 and 2 appears to be mediated by the interaction of endogenous TCF protein with its DNA target sites, as it is only observed with TOPFLASH, which contains five optimal TCF binding sites, but not with the control reporter FOPFLASH, which contains five mutated sites (Morin, Sparks et al. 1997). Thus this experiment adds supportive evidence to the notion that Doll proteins transduce Wnt signals by activating TCF target genes in a β -Catenin-dependent manner.

In summary, the protein-protein interactions demonstrated between *Drosophila* Doll and Lgs and those between their human homologues human Doll and hLgs/Bcl9, respectively, in conjunction with the genetic and cell biological data show that Doll proteins are positive regulators of the Wg and Wnt signaling pathways, respectively.

EXAMPLES**Example I: Isolation of *doll* cDNA**

The cDNA for Daughter of Legless (*Doll*) was isolated in two independent yeast genetic screens of a *Drosophila* cDNA-library for proteins directly binding to Lgs. Other DNA libraries can be used as well, such a genomic and cDNA libraries from vertebrate and invertebrate organisms. Other methods than a yeast-two hybrid screen can be used as well. Such methods include, but are not limited to, direct amplification using gene specific primers and standard methods known by people skill in the art. To perform the yeast two-hybrid screening for protein binding to Lgs cDNA sequences encoding the first 732 amino acids ("LgsN") and the full-length protein of 1464 amino acids ("LgsFL") were subcloned into a yeast expression vector (pLexA, Clontech), fusing them to the LexA DNA-binding domain. Subsequently these constructs were transformed into the *LEU2*-reporter yeast strain EGY48 together with the *lacZ*-reporter plasmid pSH18-34 and an embryonic *Drosophila melanogaster* cDNA-library fused to an acidic transcriptional activation domain ("RFLY-1" library, PNAS 93, 3011-3015). In a first step triple-transformant colonies containing the LgsN- or LgsFL-LexA-fusion constructs, respectively, the pSH18-34 reporter and a RFLY-1 library plasmid were grown on minimal selective medium plates for two days, harvested, thoroughly mixed, and stored as uniform aliquots. Then cells from one of these aliquots were transferred into permissive Galactose/Raffinose minimal selective liquid medium, and incubated with shaking at 30°C for a few hours, thereby inducing expression of the library cDNA-activation domain fusion from the *GAL1*-inducible promotor. Finally these "induced" cells were plated on Galactose/Raffinose minimal selective medium plates lacking the amino acid l-leucine. On these plates cell growth was sustained only upon activation of a *LEU2*-selector gene through molecular interaction of the respective LexA-fusion and activation domain-fusion proteins. The *LEU2*-gene codes for an essential metabolic enzyme needed for the biosynthesis of leucine from other amino

acid precursors. All clones growing under these restrictive conditions were isolated and analyzed for the activity of the *lacZ*-reporter gene, encoding the metabolic enzyme β -Galactosidase from the enterobacterium *E.coli*, by a standard X-Gal assay (e.g. Bartel and Fields (eds.), Oxford University Press 1997). From all candidate clones that passed these two selection steps, the cDNA-library plasmids were isolated again by standard techniques (e.g. Methods in Yeast Genetics, Cold Spring Harbour Laboratory Press, 1997) and retested for specific interaction with Lgs in the X-Gal assay, using an unrelated LexA-fusion protein as a negative control. By this procedure three independent cDNA-clones were identified, that strongly and specifically interacted only with Lgs and contained partially overlapping sequences: BK12b, BK14b and TK5.35h. By searching the *Drosophila* genome database using the blastn algorithm (<http://www.ncbi.nlm.nih.gov:80/BLAST/>) we mapped the three isolated cDNAs to the CG11518 locus, coding for a protein product of 815 amino acids in length. The cDNA-clones coded different parts of the Doll protein, with BK12b containing nucleotides (nt) 2223-2448, BK14b nt 2191-2448 and TK4.35h nt 749-2448 of the computationally predicted open reading frame (ORF). Further bioinformatical analysis (<http://www.ebi.ac.uk/interpro/>) revealed that the very C-terminal part of the protein sequence (ca. aa 745-805), present in all three of the Lgs-binding clones, was predicted to adapt a PHD-finger fold, which has been identified in other proteins involved in transcriptional regulation at different levels.

Example II: Identification of Human and Mouse Homologues of *Drosophila* Doll

After the identification of the *Drosophila* Doll amino acid sequence, publicly available databases were searched for similar protein sequences in other species, using the tblastn algorithm (<http://www.ncbi.nlm.nih.gov:80/BLAST/>). Two candidate sequences were found each in ESTs from *Mus musculus* and *Homo sapiens*, respectively, the putative protein products of which display high

similarity to Doll in their C-terminal domains. These stretches of high similarity are predicted to adapt a PHD-finger fold as well (Figure 4). Doll proteins do not display other known structural motifs in their N-terminal sequences but they display a second high homology domain, which was accordingly named DOLL Homology Domain (DHD) (Figure 4). Doll proteins of both invertebrate and vertebrate origin have so far not been further described or experimentally studied, and have thus not previously been implicated in any specific biological process.

Example III: Isolation and mapping of *Drosophila* doll alleles

EMS-treated males were crossed to females carrying a wg transgene (sev-wg) driven by two copies of the sevenless enhancer (Basler, Christen et al. 1991). 2×10^5 progeny were screened for suppressors of the rough eye phenotype. Third chromosomal suppressors were coarsely mapped by meiotic recombination using a panel of P[y +] insertions. One such suppressor, Sup¹³⁰, showed intriguingly dominant lethality in combination with the lgs allele lgs^{17B} (US09/915.543) (Sup¹³⁰ /+ lgs^{17B} /+ transheterozygous animals do not survive), strongly suggesting a close genetic interaction. Fine mapping of the mutation using denaturing HPLC (WAVE system, Transgenomic Inc.) demonstrated that it localizes within the doll gene. The doll coding region was therefore sequenced using PCR fragments covering the doll coding region derived from genomic DNA from homozygous Sup¹³⁰ mutant larvae. The defect in Sup¹³⁰ was found to be a 14 bp deletion (nucleotides 2253 to 2266: 5' CATGTGCCACAAGG 3') within the doll open reading frame that induced a frame-shift subsequent to amino acid 751 and resulted in the formation of a premature stop codon. Hence this allele is referred to as doll¹³⁰ and encodes a truncated Doll protein lacking the C-terminal PHD finger.

Pole cell transplantation, chromosome squashes, and chromosome in situ hybridization experiments were carried out according to standard protocols (Ashburner 1989).

Example IV: Use of doll as a hybridization probe

The following method describes the use of a non-repetitive nucleotide sequence of *doll* as a hybridization probe. The method can be applied to screen for *doll* homologues in other organisms as well. DNA comprising the sequence of *doll* (as shown in Figures 1,2,3) is employed as probe to screen for homologue DNAs (such as those included in cDNA or genomic libraries). Hybridization and washing of the filters containing either library DNAs is performed under standard high stringency conditions (Sambrook, Fritsch et al. 1989). Positive clones can be used to further screen larger cDNA library platings. Representative cDNA-clones are subsequently cloned into pBluescript (Stratagene) or similar cloning vectors and sequenced.

Example V: Use of *doll* as a hybridization probe for *in situ* hybridization.

In situ hybridization of *Drosophila doll* mRNA can be performed in embryo as described in (Tautz and Pfeifle 1989). However, with small modifications it can also be used to detect any mRNA transcript in *Drosophila* larval imaginal discs or vertebrate tissue sections. Labeled RNA probes can be prepared from linearized *doll* cDNA (as showed in Figures 1,2,3), or a fragment thereof, using the DIG RNA labeling Kit (SP6/T7) (Boehringer Mannheim) following the manufacturer's recommendations.

Example VI: Expression of *doll* in *Drosophila melanogaster*

Doll can be expressed in *Drosophila* in the whole organism, in a specific organ or in a specific cell type, during the whole life or only at a specific developmental stage, and at different levels. An overview of the standard methods used in *Drosophila* genetics can be found in (Brand and Perrimon 1993; Perrimon 1998; Perrimon 1998).

Generation of *doll* mutant embryos

Mosaic germlines are generated with the help of site-specific recombination through the FLP recombinase (Xu and Rubin 1993).

Females of the genotype *hsp70:flp*, *FRT82 doll¹³⁰* / *FRT82 ubi-GFP* are heat-shocked at 37°C for 1 hr during the third instar larval stage to induce FLP-directed recombination and later mated to *doll¹³⁰* / *TM6b[y+]* males. Germline mosaics are induced. The source of recombinase is a first chromosome insertion of a fusion of the *hsp70* promoter (denoted by "*hsp70*") to the FLP coding sequence. Somatic recombination at the *FRT82* sites gives rise to adult female germ line that produces oocytes that upon fertilization lead to embryos which do not contain neither zygotic nor maternally contributed information for the production of functional dDoll protein. Those embryos can be identified by the absence of the yellow+ phenotype provided by the *TM6b[y+]* paternal balancer chromosome. For analysis, cuticles are prepared by standard techniques from mutant embryos, and examined by dark field microscopy.

dAPC2 *doll* doublemutant germ line clones were generated with an *FRT82 dAPC2 DS doll¹³⁰* chromosome. The *FRT82 ovo^{D1}* chromosome (Chou and Perrimon 1996) was used to select for mutant germ cells. The *FRT82 doll¹³⁰* chromosome was also used to create *doll* mutant clones in discs, in conjunction with an *FRT82 arm-lacZ* chromosome.

Generation of doll mutant embryos expressing constitutively active Arm

In order to express constitutively active Arm (" Δ Arm"), females of the genotype described above are heat shocked at 37°C for 1 hr during late pupal stages and mated to males of the genotype *UAS: Δ Arm hsp70-Gal4* / *UAS: Δ Arm hsp70-Gal4; doll* / *TM6b[y+]*. Due to the presence of the additional transgenes in these males offspring that had arisen from a *doll* mutant oocytes and *doll* mutant sperm express upon heat treatment the constitutively active Arm protein, that transiently induced Wingless target genes.

Example VII: Rescue of *ddoll*-/- flies with *hdoll*-1 and *hdoll*-2 cDNA expression

In order to confirm the functional homology between *Drosophila* and human Doll-1 and human Doll-2, the human genes were introduced into *Drosophila* flies carrying two mutant *doll* alleles (*ddoll*-/- flies). Specifically, flies carrying e.g. a

tub:hdoll transgene, and two mutant doll alleles, e.g. doll¹³⁰ and EP(3)1076 (publicly available) were generated. ddoll^{-/-} mutant flies display larval or pupal lethality. In contrast, ddoll^{-/-} mutant flies carrying at least one copy of the tub:hdoll-1 or tub:hdoll-2 transgenes survive to adulthood. This demonstrates that both, hDoll-1 and hDoll-2, can replace endogenous dDoll function in flies and thus validates functional homology between *Drosophila* and human Doll (Figure 7).

Example VIII: Protein production and purification of Doll in *E. coli*

The following method describes recombinant expression of Doll proteins in bacterial cells. DNA encoding full-length or a truncated Doll form is fused e.g. downstream of an epitope tag or glutathione-S-transferase (GST) cDNA and a thrombin or enterokinase cleavage site contained within an inducible bacterial expression vector. Such epitope tags include poly-his, S-protein, thioredoxin and immunoglobulin tags. A variety of plasmids can be employed, including commercially available plasmid such as pGEX-4T (Pharmacia) or pET-32a (Novagen).

Briefly, a bacterial expression plasmid containing the doll sequence, for instance fused to a GST-sequence, is transformed by conventional methods into protease deficient *E. coli* such as BL21 (Stratagene). A bacterial colony containing the plasmid is then expanded overnight in selection medium to reach saturation. The next morning, this culture is diluted 1:100 and bacterial are allowed to growth to an optical density (OD₆₀₀) of 0.6. Protein production is initiated by addition of an inducer of the promoter under which GST-Doll fusion protein is expressed. A variety of inducers can be employed depending on the expression vector used, including IPTG.

Expressed GST tagged Doll can then be purified, for instance, using affinity beads or affinity chromatography, such as glutathione beads (commercially available from Pharmacia). Extracts are prepared by lysing the Doll-expressing bacteria in sonication buffer (10 mM Tris HCl pH 8.0, 150 mM NaCl, 1 mM EDTA, 1.5% sarkosyl, 2% Triton-X-100, 1 mM DTT and protease inhibitors), followed by short sonication on ice (e.g. 3 times 20 seconds at

middle power) and centrifugation. Cleared supernatants are then incubated under gentle rotation for example with glutathione beads for 1 hrs at 4°C. Next beads are washed several time in washing buffer (20 mM Tris pH 8.0, 200 mM NaCl, 1 mM EDTA, 1 mM DTT, 1 mM MgCl₂, 0.5% NP40), and finally stored in storage buffer (20 mM Tris pH 8.0, 200 mM NaCl, 1 mM EDTA, 1 mM DTT, 1 mM MgCl₂, 10 % glycerol, 0.5% NP40, and proteinase inhibitors).

Alternatively, a His-tagged, S-protein, thioredoxin or IgG tagged Doll can be purified using affinity chromatography.

The quality of the preparations can be checked e.g. by SDS-gel electrophoresis and silver staining or Western blot.

In case the epitope-tag has to be cleaved, several methods are available depending on the presence of a cleavage site between the epitope-tag and the Doll protein. For example, it is possible to produce a GST-Doll fusion protein containing a thrombin cleavage site right before the first Doll amino acid. Briefly, a GST-Doll preparation on glutathione-affinity beads is washed several times in cleavage buffer (50 mM Tris HCl pH 7.0, 150 mM NaCl, 1 mM EDTA, 1 mM DTT). Thrombin is then added and the samples are incubated for over 16 h at room temperature. Supernatants are then collected and analyzed for successful cleavage of Doll from the beads by polyacrylamide gel electrophoresis and silver staining or Western blot.

Example IX: Protein-protein interactions involving Doll

A GST-fusion protein in vitro binding assay can be performed to map binding domains and find additional interaction partners. For this purpose, proteins are in vitro translated using reticulocyte lysates (e.g. TNT-lysates, Promega Corporation) containing [³⁵S]methionine following the instructions provided by the manufacturer. Alternatively, cellular proteins can be labeled by incubation of culture cells with [³⁵S]methionine. Glutathione S-transferase (GST) fusion proteins, produced as illustrated in the Example VIII, are immobilized on glutathione-Sepharose and blocked in binding buffer (20 mM Tris pH 8.0, 200 mM NaCl, 1 mM EDTA, 1 mM DTT, 1 mM MgCl₂, 10 % glycerol, 0.5% NP40, 0.05% BSA,

and proteinase inhibitors) for 45 min. Two μ g of immobilized GST proteins are then incubated for 1.5 hrs with 0.5-6 μ l of in vitro translated proteins in binding buffer or with [35 S]methionine labeled cell extract. The beads are washed four times in washing buffer (20 mM Tris pH 8.0, 200 mM NaCl, 1 mM EDTA, 1 mM DTT, 1 mM $MgCl_2$, 0.5% NP40) and boiled in Laemmli SDS sample buffer. Proteins binding to Doll are detected by autoradiography. In case that a cell lysate were used to identify novel Doll binding partner, the protein bands on the gel can be isolated by methods known in the art, and the protein sequence can be determined e.g. by mass spectrophotometrical analysis.

A yeast two hybrid assay can additionally be performed to confirm the results of the in vitro binding assays described above or to screen cDNA library for new interaction partners (Fields and Sternglanz 1994). In this context, the desired cDNAs are subcloned into appropriate yeast expression vectors that link them either to a Lex DNA binding domain (e.g. pLexA, Clontech) or an acidic activation domain (e.g. pGJ4-5, Clontech). The appropriate pair of plasmids is then transformed together with a reporter plasmid (e.g. pSH18-34, Clontech) into an appropriate yeast strain (e.g. EGY48) by the lithium acetate-polyethylene glycol method and grown on selective media (Sambrook, Fritsch et al. 1989). Transformants are analyzed for reporter gene activity as described by the manufacturer of the vector-reporter plasmid used. To establish reproducibility the interactions is tested in both directions. Alternatively, this method is used to screen for novel Doll interaction partners. In this context, e.g. pLexA-Doll is transfected into yeast together with a cDNA library cloned into e.g. pGJ4-5 as described above. Positive clones can be isolated and the cDNA they contain can be sequenced by methods known by people skilled in the art.

Example X: Immunohistochemistry

Localization of the Doll proteins is performed on *Drosophila* embryo, imaginal discs, invertebrate and vertebrate adult tissue sections or tumor cell lines using the anti-Doll antibodies provided by this invention. For instance, if a tumor cell line is used, cells can be seeded into polylysine-coated 8 well chambers (Nalge-Nunc Internat.) and grown overnight at 37°C. As a positive control, 293 MEK cells (ATCC) cells might be transfected

e.g. by a lipofection method (e.g. Lipofectamine, Gibco technologies) with a Doll expression plasmid, such as pcDNA3.1 (Invitrogen). Two days after transfection, cells are washed and fixed with 3.7% formaldehyde in PBS for 10 min, permeabilized in 0.5% Triton-X-100 for another 10 min, and blocked with a 1:1000 dilution of pre-immunoserum in 2% BSA-PBS for 1h at RT. Cells are then incubated with a 1:1000 dilution of anti-Doll immunoserum for 2 hrs at RT, followed by washing in PBS and staining with anti-rabbit secondary-antibody. The washing step is repeated and preparations are blocked in a solution of 3% BSA in PBS/0.1% TritonX-100 for 1 hr. The slides are then washed three times for 5 min in PBS and incubated with a 1:200 dilution (v/v) of TRITC-conjugated swine anti-rabbit immunoglobulin (Dako, Inc.). The washing step is repeated before applying coverslips using Vectashield® mounting medium (Vector Laboratories, Inc.). Detection of other proteins such as β -Catenin, hLgs or Tcf can be performed in the same way using anti- β -Catenin (commercially available), anti-hLgs (US 09/915.543) or anti-Tcf (commercially available) specific antibodies, respectively.

Example XI: Luciferase reporter gene assays

The effect of Doll on Tcf transactivation activity can be performed in a cell culture system using a Tcf responsive luciferase reporter gene. Depending on the expression vector used, this protocol can be applied for mammalian as well as for *Drosophila* cell lines. For instance, HEK293 cells (ATCC) are a well suitable system. Hereby, Doll full length cDNA is cloned into a mammalian expression vector, such as pcDNA3 (Invitrogen), and transfected together with the TOPFLASH luciferase reporter plasmid (Upstate biotechnology, New York, USA) into 293 cells. A lipofection agent like the Lipofectamine transfection reagent (Life Technologies, Inc.) can be used for this purpose. A renilla luciferase reporter plasmid, e.g. pRL-SV40, (Promega Corporation, Madison USA), is co-transfected to normalize the transfection efficiency. Cell extracts are prepared 48 h after transfection and assayed for firefly and renilla luciferase activity as described by the manufacturer (Dual luciferase reporter assay system, Promega Corporation). All the luciferase

values are normalized for renilla luciferase activity (see Figure 8).

Example XII: Screening of chemical compounds, organic products or peptides interfering with Doll function

A reporter gene assay is performed with a similar protocol as described in example XI, but scaled down to be performed as a high throughput screening. For this purpose colon cancer cell lines with mutated and/or constitutively active β -Catenin are stably transfected with the Topflash vector described in Example XI and Doll cDNA. The established monoclonal population, which gives the most reliable and constant reporter gene activity is selected for later assays. One day after plating, cells are treated with single compounds derived from a chemical or peptide library. One to 24 hours later reporter gene activity is measured. Compounds found to inhibit reporter gene activity are then further characterized for specific activity on the Doll-containing transcriptional complex. Alternatively, Wnt pathway activity can be measured by detecting mRNA or protein levels of a target gene, e.g. myc (He, Sparks et al. 1998).

Example XIII: Screening assay based on protein-protein interaction for compounds inhibiting Doll-Lgs or Doll interaction partner X

Doll and its interaction partner or fragments thereof are produced and purified e.g. from E.coli cultures (e.g. as described in example VIII). Proteins are tagged e.g. with 6 histidines, S-protein, GST or thioredoxin. Small aliquots of the purified proteins are incubated in an appropriate binding buffer. At this point chemical compounds are added to the mixture and their capacity to disrupt the protein-protein interaction is monitored e.g. by any of the methods described below. Compounds inhibiting this interaction are subsequently tested for their specificity and in vivo toxicity. Well established methods to monitor protein-protein interactions are e.g.:

- Time resolved fluorometry with lanthanide chelate labels (Hemmilä I. And Webb S. DDT 2: 373-381 (1997))
- Scintillation proximity assay (SPA) (Amersham life Science)

- Fluorescence polarisation

References

- Aasland, R., T. J. Gibson, et al. (1995). "The PHD finger: implications for chromatin-mediated transcriptional regulation." Trends Biochem Sci 20(2): 56-9.
- Ashburner, M. (1989). Drosophila A laboratory handbook. Cold Spring Harbor, New York, Cold Spring Harbor Laboratory Press.
- Baker, N. E. (1988). "Transcription of the segment-polarity gene wingless in the imaginal discs of *Drosophila*, and the phenotype of a pupal-lethal wg mutation." Development 102(3): 489-97.
- Barker N, H. G., Korinek V, Clevers H (1999). "Restricted high level expression of Tcf-4 protein in intestinal and mammary gland epithelium." Am J Pathol 154: 29-35.
- Basler, K., B. Christen, et al. (1991). "Ligand-independent activation of the sevenless receptor tyrosine kinase changes the fate of cells in the developing *Drosophila* eye." Cell 64(6): 1069-81.
- Brand, A. H. and N. Perrimon (1993). "Targeted gene expression as a means of altering cell fates and generating dominant phenotypes." Development 118(2): 401-15.
- Cabrera, C. V., M. C. Alonso, et al. (1987). "Phenocopies induced with antisense RNA identify the wingless gene." Cell 50(4): 659-63.
- Chou, T. B. and N. Perrimon (1996). "The autosomal FLP-DFS technique for generating germline mosaics in *Drosophila melanogaster*." Genetics 144(4): 1673-9.
- Fields, S. and R. Sternglanz (1994). "The two-hybrid system: an assay for protein-protein interactions." Trends Genet 10(8): 286-92.
- Grosschedl R, E. Q. (1999). "Regulation of LEF-1/TCF transcription factors by Wnt and other signals." Current Opinion in Cell Biology 11: 233-240.

- He, T. C., A. B. Sparks, et al. (1998). "Identification of c-MYC as a target of the APC pathway [see comments]." Science 281(5382): 1509-12.
- Heberlein, U., E. R. Borod, et al. (1998). "Dorsoventral patterning in the Drosophila retina by wingless." Development 125(4): 567-77.
- Morin, P. J. (1999). "beta-catenin signaling and cancer." Bioessays 21(12): 1021-30.
- Morin, P. J., A. B. Sparks, et al. (1997). "Activation of beta-catenin-Tcf signaling in colon cancer by mutations in beta-catenin or APC [see comments]." Science 275(5307): 1787-90.
- Nusse, R. and H. E. Varmus (1982). "Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome." Cell 31(1): 99-109.
- Peifer, M. and P. Polakis (2000). "Wnt signaling in oncogenesis and embryogenesis--a look outside the nucleus." Science 287(5458): 1606-9.
- Perrimon, N. (1998). "Creating mosaics in Drosophila." Int J Dev Biol 42(3): 243-7.
- Perrimon, N. (1998). "New advances in Drosophila provide opportunities to study gene functions." Proc Natl Acad Sci U S A 95(17): 9716-7.
- Perrimon, N. and A. P. Mahowald (1987). "Multiple functions of segment polarity genes in Drosophila." Dev Biol 119(2): 587-600.
- Polakis, P., M. Hart, et al. (1999). "Defects in the regulation of beta-catenin in colorectal cancer." Adv Exp Med Biol 470: 23-32.
- Potter, J. D. (1999). "Colorectal cancer: molecules and populations." Journal of the National Cancer Institute 91(11): 916-32.
- Rijsewijk, F., M. Schuermann, et al. (1987). "The Drosophila homolog of the mouse mammary oncogene int-1 is identical to the segment polarity gene wingless." Cell 50(4): 649-57.

- Roose, J. and H. Clevers (1999). "TCF transcription factors: molecular switches in carcinogenesis." Biochimica et Biophysica Acta 1424(2-3): M23-37.
- Sambrook, J., E. F. Fritsch, et al. (1989). Molecular cloning: a laboratory manual, Cold Spring Harbor Laboratory Press.
- Sharma, R. P. and V. L. Chopra (1976). "Effect of the Wingless (wgl) mutation on wing and haltere development in *Drosophila melanogaster*." Dev Biol 48(2): 461-5.
- Struhl, G. and K. Basler (1993). "Organizing activity of wingless protein in *Drosophila*." Cell 72(4): 527-40.
- Tatusova TA, M. T. (1999). "Blast 2 sequences - a new tool for comparing protein and nucleotide sequences." FEMS Microbiol Lett. 174: 247-250.
- Tautz, D. and C. Pfeifle (1989). "A non-radioactive in situ hybridization method for the localization of specific RNAs in *Drosophila* embryos reveals translational control of the segmentation gene hunchback." Chromosoma 98(2): 81-5.
- van de Wetering, M., R. Cavallo, et al. (1997). "Armadillo coactivates transcription driven by the product of the *Drosophila* segment polarity gene dTCF." Cell 88(6): 789-99.
- Waltzer, L. and M. Bienz (1999). "The control of beta-catenin and TCF during embryonic development and cancer." Cancer & Metastasis Reviews 18(2): 231-46.
- Willis, T. G., I. R. Zalcberg, et al. (1998). "Molecular cloning of translocation t(1;14)(q21;q32) defines a novel gene (BCL9) at chromosome 1q21." Blood 91(6): 1873-81.
- Wodarz, A. and R. Nusse (1998). "Mechanisms of Wnt signaling in development." Annual Review of Cell & Developmental Biology 14: 59-88.
- Xu, T. and G. M. Rubin (1993). "Analysis of genetic mosaics in developing and adult *Drosophila* tissues." Development 117(4): 1223-37.

Claims

1. A nucleic acid sequence coding for a polypeptide being part of at least one signaling pathway in insects and vertebrates, characterized in that said nucleotide sequence is the "Daughter of Legless" (DOLL) gene as well as homologues, fragments, derivatives and functional and structural analogs thereof.
2. The nucleic acid sequence according to claim 1, characterized in that said signaling pathway is the Wnt signaling pathway.
3. The nucleic acid sequence according to claim 2, characterized in that it is the *Drosophila melanogaster* *doll* gene (*ddoll*) comprising the nucleotide sequence as shown in SEQ. ID. NO 1.
4. The nucleic acid sequence according to claim 3, characterized in that it is coding for a polypeptide comprising 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the *Drosophila* Doll polypeptide as shown in SEQ. ID. NO 6.
5. The nucleic acid sequence according to claim 2, characterized in that it is the human *doll-1* (*hdoll-1*) comprising the nucleotide sequence as shown in SEQ ID. NO 2.
6. The nucleic acid sequence according to claim 5, characterized in that it is coding for a polypeptide comprising 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the human Doll (*hDoll-1*) polypeptide as shown in SEQ. ID. NO 7.

7. The nucleic acid sequence according to claim 2, characterized in that it is the human *doll-2* gene (*hdoll-2*) having the nucleotide sequence as shown in SEQ ID. NO 3.
8. The nucleic acid sequence according to claim 7, characterized in that it is coding for a polypeptide comprising 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the human Doll polypeptide (*hDoll-2*) as shown in SEQ. ID. NO 8.
9. The nucleic acid sequence according to claim 2, characterized in that it is the mouse *doll-1* gene (*mdoll-1*) comprising the nucleotide sequence as shown in SEQ ID. NO 4.
10. The nucleic acid sequence according to claim 9, characterized in that it is coding for a polypeptide comprising 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the mouse Doll (*mDoll-1*) polypeptide as shown in SEQ. ID. NO 9.
11. The nucleic acid sequence according to claim 2, characterized in that it is the mouse *doll-2* gene (*mdoll-2*) comprising the nucleotide sequence as shown in SEQ ID. NO 5.
12. The nucleic acid sequence according to claim 11, characterized in that it is coding for a polypeptide comprising 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the mouse Doll polypeptide (*mDoll-2*) as shown in SEQ. ID. NO 10.
13. A polypeptide being part of at least one signaling pathway in insects and vertebrates, characterized in that said polypeptide is the "Daughter of Legless" (DOLL)

protein as well as homologues, fragments, derivatives and structural and functional analogs thereof.

14. The polypeptide according to claim 13, characterized in that said signaling pathway is the Wnt signaling pathway.
15. The polypeptide according to claim 14, characterized in that said polypeptide is the *Drosophila melanogaster* Doll protein having the amino acid sequence as shown in SEQ. ID. NO. 6.
16. The polypeptide according to claim 15, characterized in that said polypeptide has a 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the Doll polypeptide as shown in SEQ. ID. NO 6.
17. The polypeptide according to claim 14, characterized in that said polypeptide is the human Doll protein (hDOLL-1) having the amino acid sequence as shown in SEQ. ID. NO. 7.
18. The polypeptide according to claim 17, characterized in that said polypeptide has a 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the Doll polypeptide as shown in SEQ. ID. NO 7.
19. The polypeptide according to claim 14, characterized in that said polypeptide is the human Doll protein (hDoll-2) having the amino acid sequence as shown in SEQ. ID. NO. 8.
20. The polypeptide according to claim 19, characterized in that said polypeptide has a 50% to 100%, preferably 100%

sequence identity to a fragment or the entire sequence of the Doll polypeptide as shown in SEQ. ID. NO 8.

21. The polypeptide according to claim 14, characterized in that said polypeptide is the mouse Doll-1 protein (mDoll-1) having the amino acid sequence as shown in SEQ. ID. NO. 9.
22. The polypeptide according to claim 21, characterized in that said polypeptide has a 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the Doll polypeptide as shown in SEQ. ID. NO 9.
23. The polypeptide according to claim 14, characterized in that said polypeptide is the mouse Doll-2 protein (mDoll-2) having the amino acid sequence as shown in SEQ. ID. 10.
24. The polypeptide according to claim 23, characterized in that said polypeptide has a 50% to 100%, preferably 100% sequence identity to a fragment or the entire sequence of the Doll polypeptide as shown in SEQ. ID. NO 10.
25. The nucleic acid sequence according to claim 2, characterized in that said nucleic acid sequence is coding for a polypeptide comprising a low overall amino acid sequence identity and a sequence identity of 50% to 100%, preferably 100% in conserved domains.
26. The nucleic acid sequence according to claim 25, characterized in that said conserved domain is the DHD domain.
27. The nucleic acid sequence according to claim 26, characterized in that said conserved domain is the PHD finger.
28. The nucleic acid sequence according to claim 2, characterized in that said sequence comprises a fragment of 20

to 100 nucleotides in length, preferably from 20 to 60 nucleotides, and most preferably from 20 to 50 nucleotides, said fragment being derived from SEQ. ID. NO. 1, 2, 3, 4 or 5.

29. The nucleic acid sequence according to claim 28, used as a hybridization probe.
30. A vector comprising a nucleic acid molecule encoding a *Drosophila* or vertebrate *doll* gene or a fragment thereof, selected from the group consisting of SEQ. ID. NO. 1, 2, 3, 4 or 5.
31. The vector according to claim 30, selected from the group consisting of eucaryotic and procaryotic expression vectors.
32. A host cell comprising the vector of claim 31, selected from the group consisting of mammalian cells, yeast cells, plant cells, insect cells or bacterial cells.
33. A method for the preparation of Doll proteins, fragments, derivatives and analogs thereof, comprising the steps of:
 - a) isolating a nucleic acid sequence containing the complete *doll* cDNA sequence or part thereof;
 - b) recombinantly expressing *doll* cDNA or a fragment thereof in bacterial, mammalian, plant, yeast or insect cells;
 - c) inducing protein production in said cells;
 - d) purifying Doll proteins.
34. The method according to claim 33, wherein step a) comprises the isolation of a nucleic acid molecule encompassing the *doll* cDNA from a vertebrate or invertebrate cDNA or genomic library.

35. The method according to claim 33, wherein step b) comprises the fusion of DNA encoding full-length or truncated doll to an epitope tag and a cleavage site contained within an inducible eucaryotic or procaryotic expression vector and transforming the appropriated host cells with said expression vector.
36. A chimeric protein comprising a Doll polypeptide fused to a heterologous amino acid sequence selected from the group consisting of an epitope-tagged sequence, an antibody, glutathione-S-transferase protein, β -galactosidase, and alkaline phosphatase.
37. The chimeric protein according to claim 36, characterized in that the Doll polypeptide is selected from the group consisting of dDoll, hDoll-1, hDoll-2, mDoll-1 and mDoll-2.
38. The polypeptide according to claim 14, characterized in that it comprises the full length Doll polypeptide or a fragment thereof comprising an antibody-binding site for an anti-Doll antibody.
39. An antibody specifically recognizing a Doll polypeptide, said antibody being selected from the group consisting of polyclonal and monoclonal antibodies and fragments thereof.
40. An assay for studying diseases induced by a disrupted Wnt function or for drug screening comprising the use of organisms selected from the group consisting of Drosophila, mice, rats, rabbits, chicken, frogs, pigs or sheep, said organisms showing increased or reduced or no expression of doll or express a mutated Doll polypeptide in at least one tissue or organ.

41. The assay according to claim 40, characterized in that said organisms express the *doll* gene as a heterologous transgene.
41. The assay according to claim 40, characterized in that said *doll* gene comprises a mutation selected from the group consisting of deletions, point mutations, foreign DNA insertions and inversions.
42. Use of a Doll protein, homologue, derivative and fragment thereof for the development of a therapeutic and diagnostic method for the treatment of conditions selected from the group consisting of disorders of cell fate, differentiation or proliferation.
43. Use of a Doll protein, homologue, derivative and fragment thereof for the development of a therapeutic and diagnostic compound for the diagnosis of conditions selected from the group consisting of disorders of cell fate, differentiation or proliferation.
44. The use of a Doll protein, homologue, derivative and fragment thereof according to claim 43, characterized in that said therapeutic or diagnostic compound is selected from the group consisting of *Drosophila* and vertebrate Doll protein homologues and fragments thereof and antibodies and antibody fragments thereof.
45. Use of *doll* nucleic acids, homologues, derivatives and fragments thereof for the development of a therapeutic and/or diagnostic compound for the treatment of conditions selected from the group consisting of disorders of cell fate, differentiation or proliferation and its application to an individual.

46. Use of doll nucleic acids, homologues, derivatives and fragments thereof for the development of a therapeutic and diagnostic method for the treatment of conditions selected from the group consisting of disorders of cell fate, differentiation or proliferation and its application to an individual.
47. The use of doll nucleic acids, homologues, derivatives and fragments thereof according to claim 45, characterized in that said therapeutic or diagnostic compound is selected from the group consisting of doll antisense DNA or RNA, doll double-stranded RNA and chemical or natural occurring compounds interfering with doll function.
48. A peptide comprising a fragment of the Doll polypeptide.
49. The peptide according to claim 48, comprising 40 to 60 amino acids in the N-terminal region of hDoll-1, hDoll-2, mDoll-1, mDoll-2 and Drosophila Doll.
50. The peptide according to claim 49, said peptide comprising the Doll homology domain (DHD).
51. Use of DHD in screening methods for the identification of chemical compounds, organic products, polypeptides or peptides interfering with Doll function in the Wnt pathway.
52. The use of DHD according to claim 51 in screening assay based on protein-protein interactions.
53. The use of DHD according to claim 52, wherein said screening assay is selected from the group comprising an *in vitro* protein-protein interaction assay and a protein-protein interaction assay in a host cell.

54. The use of DHD according to claim 51, in a screening assay for compounds specifically inhibiting the interaction between Doll and another protein.

Figure 1: The translated sequence of the *Drosophila* doll gene.

dDoll cDNA Sequence Range: 1 to 2448

ATGACCCACAATCTTGGTATGGCGCCATATCGATTGCCGGGTCCAGCGGG
CGGACTCTGTCCGCCCCGATTTTAAGCCGCCGCTCCCACGGACATCATCT
CGGCGCCGAGCAATCCGAAGAAGCGGCGAAAAACCTCAAGTGCCGCCAAC
TCCGCTGCAGCGGTGGCTGCGGCGGCGGCTGCAGCAGCTGCTGCGAATTC
CATGCAGCAGCAGCAGGCGCCACCCACACCGCAGGATTTGCTGCCCCCTC
CGCCAATGGGAGGCTTCGGAGACACCATATTGCCTCGAATCCATTTCGAC
GACAGTCCCCAGGTGTCTGGCGATGTCCAGCTCAGCGGCCGCGGCGATGGC
GGCCATGAATCAGATGGGCGGCGGACCAGGAGGTGGTCACTTTGGCGGCG
GTGGACCGGGTGGGCACCCGCACTGGGAAGACCGCATGGGCATGGGCGGT
GGACCTCCTCCCCCGCCTCACATGCATCCCCATATGCACCCGCATCATCC
AGGCGGACCTATGGGTACCCACATGGCCCACATCCGCACATGGGTGGTC
CACCTCCAATGCGAGGAATGAGCCCCATGCACCCCCATCAAATGGGACCG
GGACCAGGCGTCTGGACTACCGCCGCATATGAATCACGGAAGGCCAGGGGG
ACCTGGTGGTCTTGGAGGACCCGTCCCAATGGGTAGTCCCATGGGTGGAA
TAGCTGGCATGGGCGGCATGAGCCCAATGGGCGGAATGGGAGGCCCCAGC
ATATCACCCCATCACATGGGCATGGGTGGTCTGTCTGCCCATGGGAGGCGG
TCCCAACGGACCCAATCCGCGAGCCATGCAGGGTTCACCGATGGGCGGTCT
CGGGGCAGAACTCGCCAATGAACTCACTGCCTATGGGTTCGCCAATGGGC
AATCCAATTGGCAGCCCGTTGGGCCCTCCCTCGGGACCGGGCCCTGGGAA
TCCCGGCAATAACGGGCGGACCACAGCAGCAACAACAACCTCCGCAGC
CACCGATGAACAACGGGCAGATGGGTCTCTCTCTCTGCACAGTCCGCTC
GGAAACGGACCAACGGGTCATGGCAGTCACATGCCTGGAGGACCAATCCC
AGGACCAGGTCTTGGGCCTGGCGGCCTAGTAGGTCCCGGTGGCATCTCCC
CCGCGCACGGCAATAAECGGGGTGGTTCTGGGAACAACATGCTCGGCGGG
AATCCCGGCGGCGGCAACAGCAACAACAACGGAAGCAATACAAGTAACGC
CAGCAACAACAATCAAAATCCTCACCTCTCGCCAGCAGCCGGACGCCTGG
GAGTGCCGACGTCGATGCAGTCGAATGGACCTTCGGTATCATCGGTAGCC
TCCTCATCGGTTCCCTCGCCCGCCACGCCCACGCTCACGCCCACATCGAC
GGCCACGTCCATGTCCACGTCACTGCCTACATCCTCGCCAGCGCCGCCCG
CCATGTACCCGCATCACTCGCTAAACAGCGCCGGCCCGAGTCCGGGCATG
CCCAACTCGGGACCCAGCCCGCTGCAGTCACCAGCCGGACCCAATGGCCC
CAATAACAACAACAGCAATAACAACAACGGACCCATGATGGGCCAGATGA
TCCCGAACGCAGTTCTTATGCAGCACCAGCAGCACATGGGCGGCGGCCCA
CCTGGCCACGGGCCCCGGACCAATGCCCAGGAATGGGCATGAACCAATGCT
GCCACCGCAGCAACCCTCCCATCTTGGTCCCCCGCATCCGAATATGATGA

ACCACCCGCATCATCCGCACCATCATCCTGGCGGACCACCGCCGCACATG
ATGGGTGGACCCGGAATGCACGGCGGTCCTGCTGGAATGCCTCCTCATAT
GGGCGGAGGACCTAATCCGCACATGATGGGCGGTCCGCACGGGAACGCGG
GTCCGCACATGGGCCACGGCCACATGGGTGGAGTACCAGGTCCAGGACCC
GGACCCGGCGGCATGAACGGACCCCCGCATCCGCACATGTCCCCGCACCA
CGGACATCCGCATCACCACCACAATCCGATGGGCGGCCAGGTCCAAATA
TGTTCCGGCGGTGGTGGAGGAGGTCCCATGGGTCCCAGGTGGACCGATGGGC
AACATGGGGCCCATGGGAGGTGGCCCGATGGGCGGCCCTATGGGCGTAGG
TCCCAAGCCGATGACAATGGGCGGCGGGAAGATGTACCCGCCGGGACAGC
CAATGGTCTTTAATCCGCAGAACCCGAATGCGCCGCCCATATATCCTTGT
GGCATGTGCCACAAGGAGGTGAACGACAACGACGAAGCCGTGTTCTGTGA
ATCCGGTTGTAACTTTTTCTTTCACAGAACCTGTGTTGGCCTGACAGAGG
CGGCCTTCCAAATGCTCAACAAGGAGGTGTTTGCCGAGTGGTGCTGCGAC
AAGTGCGTGTCTTCCAAGCATATTCCCATGGTCAAGTTCAAGTGTTGA

dDoll Protein sequence 1-815

MTHNLGMAPYRLPGPAGGLCPPDFKPPPPTDIIISAPSNPKRRKTSSAAN
SAAAVAAAAAANSMQQQQAPPTPQDLLPPPPMGGFGDTIIASNPFD
DSPQVSAMSSSAAAAMAAMNQMGGGPGGGHFGGGGPGGHPHWEDRMGMGG
GPPPPPHMHPHMHPHHPGGPMGHPHGPHPHMGGPPPMRGMSPMHPHQMGF
GPGVGLPPHMHGRPGGPGGPVPMPGSPMGGIAGMGGMSPMGGMGGPS
ISPHHMGMGGLSPMGGGPNGPNPRAMQGS PMGGPGQNSPMNSLPMGSPMG
NPIGSPLGPPSGPGPGNPGNTGGPQQQQQPPQPPMNNGQMGPPLHSPL
GNGPTGHGSHMPGGPIPGPGPGPGGLVGPGGISPAHGNNPGGSGNNMLGG
NPGGGNSNNNGSNTSNASNNNQNPHLSPAAGRLGVPTSMQSNGPSVSSVA
SSSVPSPATPTLTPTSTATSMSTSVPTSSPAPPAMSPHHSLSAGPSPGM
PNSGPSPLQSPAGPNPNNNNNNNNNGPMMGQMI PNAVPMQHQQHMGGGP
PGHGPGPMPGMGMNQMLPPQQPSHLGPPHPNMNMNHPHHPHHHPGGPPPHM
MGGPGMHGGPAGMPPHMGGGPNPHMMGGPHGNAGPHMGGHGHMGGVPGPGP
GPGGMNGPPHPHMSPHHGHPHHHHNPMGGPGPNMFGGGGGGPMGPGGPMG
NMGPMMGGGPMGGPMGVGPKPMTMGGGKMYPPGQPMVFNPQNP NAPPIYPC
GMCHKEVNDNDEAVFCESGCNFFFHRTCVGLTEAAFQMLNKEVFAEWCCD
KCVSSKHIPMVKFKC

hDoll-1 cDNA sequence

hDoll-1 Protein sequence

MPAENSPAPAYKVSSHGGDSGLDGLGGPGVQLGSPDKKKRKANTQGSPFP
PLSEYAPPPNPNSDHLVAANPFDDNYNTISYKPLPSSNPYLGPGYPGFGG
YSTFRMPPHVPPRMSSPYCGPYSLRNQPHFFPQNPLGMGFNRPHAFNFGP
HDNSSFGNPSYNNALSQNVNMPNQHFRQNPANFNSQIIPPQNASQVSNPDL
ASNFVPGNNSNFTSPLESNHSFI PPPNTFGQAKAPPPKQDFTQGATKNTN
QNSSAHPPHLNMDDTVNQSNIELKNVNRNNAVNQENSRSSTEATNNNPA
NGTQNKPRQPRGAADACTTEKSNKSSLHPNRHGHSSSDPVYPCGICTNEV
NDDQDAILCEASCQKWFHRICTGMTETAYGLLTAEASAVWGCDDTQMADKD
VOLMRTRETFGPSAVGSDA

hDoll-2 cDNA sequence

CCCGGGTCCCCCACTCC**ATGG**CCGCCTCGGCGCCGCCCCCACC GGACAAG
CTGGAGGGAGGTGGCGGCCCCGCACCGCCCCCTGCGCCGCCCAGCACCGG
GAGGAAGCAGGGCAAGGCCGGTCTGCAAATGAAGAGTCCAGAAAAGAAGC
GAAGGAAGTCAAATACTCAGGGGCCCTGCATACTCACATCTGACGGAGTTT
GCACCACCCCCAACTCCCATGGTGGATCACCTGGTTGCATCCAACCCTTT
TGAAGATGACTTCGGAGCCCCCAAAGTGGGGGTTGCAGCCCCCTCCATTCC
TTGGCAGTCCTGTGCCCTTCGGAGGCTTCCGTGTGCAGGGGGGGCATGGCG
GGCCAGGTACCCCCAGGCTACAGCACTGGAGGTGGAGGGGGCCCCCAGCC
ACTCCGTCGACAGCCACCCCCCTTCCCTCCCAATCCTATGGGCCCTGCTT
TCAACATGCCCCCCCCAGGGTCTTGGCTACCCACCCCCAGGCAACATGAAC
TTTCCCAGCCAACCCTTCAACCAGCCTCTGGGTCAAACTTTAGTCCTCC
CAGTGGGCAGATGATGCCGGGGCCAGTGGGGGGATTGGTCCCATGATCT
CACCCACCATGGGACAGCCTCCAGAGCAGAGCTGGGCCACCTTCTCTG
TCCCAACGATTTGCTCAGCCAGGGGCTCCTTTTGGCCCTTCTCCTCTCCA
GAGACCTGGTCAGGGGCTCCCCAGCCTGCCGCCTAACACAAGTCCCTTTC
CTGGTCCGGACCCTGGCTTTCTTGGCCCTGGTGGTGAGGATGGGGGGAAG
CCCTTGAATCCACCTGCTTCTACTGCTTTTCCCAGGAGCCCCACTCAGG
CTCCCCGGCTGCTGCTGTTAATGGGAACCAGCCCAGTTTCCCCCCGAACA
GCAGTGGGCGGGGTGGGGGCACTCCAGATGCCAACAGCTTGGCACCCCCCT
GGCAAGGCAGGTGGGGGCTCCGGGGCCCCAGCCTCCCCCAGGCTTGGTGTA
CCCATGTGGTGCCTGTCGGAGTGAGGTGAACGATGACCAGGATGCCATTC
TGTGTGAGGCCTCCTGCCAGAAATGGTTCCACCGTGAGTGCACAGGCATG
ACTGAGAGCGCCTATGGGCTGCTGACCACTGAAGCTTCTGCCGTCTGGGC
CTGCGATCTCTGCCTCAAGACCAAGGAGATCCAGTCTGTCTACATCCGTG
AGGGCATGGGGCAGCTGGTGGCTGCTAACGATGGGTGACGCTGGTGAAGT
GGCCCAGGGAAGTGCACATGTCTCTCCCTGCTCTTCCAGGGTGATTTTTT
TGATGTTTGGCTCTTGGTCCTTGTTTCCACTGGCTTTCATCCCCATGGG
GCAGAAACAGTGGCTCCTGGGAGCAGAAAAGGAATTGAGGTGGGCAGGCA
GAAGAGCCTGGATTGCTCACTGTTTTGGGAACTTACATGTTGAGATCT

hDoll-2 protein sequence

MAASA PPPDKLEGGGGPAPPAPPPSTGRKQ GKAGLQMKSP EKKRRKSNT
QGPAYSHLTEFAPPPTPMVDHLVASNPFEDDFGAPKVGVAAPPFLGSPVP
FGGFRVQGGMAGQVPPGYSTGGGGGPQLRRQPPFPNPMGPAFNMPPQ
GPGYP PPGNMNFP SQPFNQPLGQNFSPPSGQMMPGPVGGFGPMISPTMGQ
PPRAELGPPSLSQRFAQPGAPFGPSPLQRPGLPSLPNTSPFP GPDPG
FPGPGGEDGGKPLNPPASTAFPQEPHSGSPAAAVNGNQPSFPNSSGRGG
GTPDANSLAPP GKAGGGSGPQP PGLVYPCGACRSEVNDDQDAILCEASC
QKWFHRECTGMTESAYGLLTTEASAVWACDLCLKTKEIQSVYIREGMGQL
VAANDG

Figure 3**mDoll-1 cDNA sequence**

ATGTCAGCGGAACAGGACAAGGAGCCCATCGCGCTGAAGAGAGTTAGAGG
TGGTGACAGTGGACTGGATGGGTTAGGAGGGCCCAATATACAACCTCGGAA
GCCAGATAAGAAAAAACGCAAGGCCAACACACAGGGATCTTCCTTTCTCCT
CCGCTGTCAGAGTACGCGCCACCACCGAATCCAAACTCCGACCATCTAGT
GGCTGCCAATCCGTTCGATGACAGCTACAACACCATTTCCTATAAACCAC
TGCCTTCATCTAATCCATATCTTGGCCCTGGGTATCCTGGCTTTGGAGGC
TACAGCACATTGAGAATGCCACCCACAGTCCCTCCAAGAATGTCTTCTCC
CTACTGTGGTCCTTACTCACTCAGGAATCAGCCCCACCCATTTCTCTCAGA
ATCCGTTGGGCATGGGCTTTAACCGGCCTCATGCTTTTAACTTTGGGCCG
CATGATAATTCGAATTTTGGAAACCCACCTTATAATAATGTACTGACTCA
GGACATTAACATGCCCCGTCAGCATTTTGTAGACAAGGCTCTGCTGAAAAC
TCAGTCAGATTCCCCCGCAGAATGTTGGCCAAGTGTCTAACCTGACCTC
GCATCTAATTTTGGCCCTGGGAATAATTCAAATTTTACCTCTCCGTTAGA
AACGAATCATTCGTTTATTCCACCCCCAAACGCGTTTGGCCAAGCAAAAG
CTCCACTTCCCAAACAAGACTTCACTCAAGGGGCAACCAAACCCCGAAT
CAGAATTCGTCCACTCACCCACCTCACCTAAATATGGAGGATCCAGTCAA
TCAGAGTAACGTCGAGTTAAAAAATGTCAACAGAAACAACGTTGTCCAAG
AGAACAGCCGTTTCGGGCAGCGCAGAGGCCACCAACAACCATGCGAATGGG
ACCCAGAACAAAGCCCCGGCAGCCAGGGGCGCAGCTGACCTGTGCACCCC
CGACAAAAGCCGCAAGTTCTCCCTGCTCCCCAGCCGGCATGGCCATTCTC
CCTCTGACCCTGTGTACCCGTGCGGGATTTGTACAAATGAAGTGAATGAC
GATCAGGACGCCATTCTGTGTGAAGCCTCTTGTGAGAAGTGGTTTCATCG
CATCTGCACTGGAATGACCGAAACAGCCTACGGGCTCCTGACAGCGGAAG
CATCCGCAGTGTGGGGCTGTGACACGTGCATGGCTGACAAGGATGTCCAG
CTCATGCGCACTAGAGAGGCCTTTGGTCCACCTGCCGTGGGCGGCGATGC
CTAA

mDoll-1 protein sequence

MSAEQDKEPIALKRVRGGDSGLDGLGGPNIQLGSPDKKKRKANTQGSSFP
PLSEYAPPPNPNSDHLVAANPFDDSYNTISYKPLPSSNPYLGPGYPGF
YSTFRMPPHVPPRMSSPYCGPYSLRNQPHFPQNPPLGMGFNRPHAFNFGP
HDNSNFGNPPYNNVLTQDINMPGQHFROGSAENFSQIPQNVGQVSNPDL
ASNFAFGNNSNFTSPLETNHSFIPPPNAFGQAKAPLPKQDFTQGATKTPN
QNSSTHPPHLNMEDPVNQSNVELKNVNRNNVVQENSRSLSAEATNNHANG
TQNKPRQPRGAADLCTPDKSRKFSLLPSRHGHSSSDPVYPCGICTNEVND
DQDAILCEASCQKWFHRICTGMTETAYGLLTAEASAVWGCDTCMADKDVQ
LMRTREAFGPPAVGGDA


Figure 3**mDoll-2 cDNA sequence**

ATGGCCGCCTCGGCGCCGCCCCACCGGACAAGCTGGAGGGAGGCAGCGG
CCCCGCACCGCCCCCGCGCCGCCAGCACCGGGAGGAAGCAGGGCAAGG
CCGGTCTGCAAATGAAGAGCCCTGAAAAGAAGCGAAGAAAGTCCAATACT
CAGGGTCCTGCATATTCACATCTGACGGAGTTCGCCCCACCCCGACCCC
CATGGTGGATCATCTGGTCGCTTCTAACCCTTTTGAGGATGATTTTCGGAG
CCCCTAAAGTGGGGGGCGCAGGCCCTCCGTTCTCGGCAGTCCGGTGCCC
TTTGAGAGGCTTTTCGTGTACAGGGGGGCATGGCAGGCCAGGTACCCCAAG
CTACGGCACTGGAGGAGGAGGGGGTCCCCAGCCACTTCGTGCGCAGCCCC
CTCCTTTTCCCCCAGCCCTATGGGTCCAGCTTTTAATATGCCCCCTCAG
GGTCCCTGGGGTACCCCGCCCCCTGGCAACATGAACTTTCCAGTCAACC
CTTCAACCAGTCTCTGGGCCAAAACCTTTAGCCACCTGGTGGGCAGGTGA
TGCCAGGCCCAGTAGGCGGATTTGGTCCCATGATCTCACCGACCATGGGA
CAGCCTCCTAGAGGGGAGCTGGGTCTCCTCCTCTCCCCAACGCTTTAC
CCAACCAGGAGCACCTTATGGTCCTTCTCTTCAGAGACCTGGTCAGGGAC
TCACCCAGCTGCCCTCCAACACAAGTCCCTTCCCTGGTCCAGACCCTGGT
TTTCCTGGACCTGGCGGTGAGGATGGTGGGAAGCCCTTGAACCCACCGGC
TCCCACCGCCTTTCCCCAGGAAGCACCATTCGGGCTCCCCGCTGCTGCTG
TCAATGGGAATCAGCCCAGTTTCCCCCCTAGCAGCAGTGGTCGAGGTGGG
GGCACTCCAGATGCCAACAGTCTGGCACCCCCCGGCAAGGCAGGGGGAGG
CTCAGGGCCCCAGCCTCCCCAGGCCTGGTGTACCCCTGCGGTGCCTGCC
GTAGTGAGGTAAATGATGACCAGGATGCCATTCTGTGTGAGGCCTCCTGC
CAGAAGTGTTTTACCGCGAGTGCACCGGCATGACCGAGAGTGCCTACGG
CCTGCTGACCACCGAGGCCTCTGCCGTCTGGGCCTGTGATCTTTGCCTCA
AGACCAAGGAGATCCAGTCTGTCTACATCCGAGAGGGCATGGGCCAGTTG
GTGGCTGCTAACGATGGGTGA


mDoll-2 protein sequence


MAASAPPPDKLEGGSGPAPPPAPPSTGRKQGKAGLQMKSPKKRRKSNT
QGPAYSHLTEFAPPPTPMVDHLVASNPFEDDFGAPKVGGAGPPFLGSPVP
FGGFRVQGGMAGQVPPSYGTGGGGGPQPLRRQPPFPSPMPGPAFNMPPQ
GPWGTPPPGNMNFPSQPFNQSLGQNFSPGGQVMPGPVGGFGPMISPTMG
QPPRGELGPPPLPQRFTQPGAPYGPSLQRPQGGLTQLPSNTSPFPGPDGP
FPGPGGEDGGKPLNPPAPTAFPQEAFFGLPAAAVNGNQPSFPPSSSGRGG
GTPDANSLAPP GKAGGGSGPQPPGLVYPCGACRSEVNDDQDAI LCEASC
QKWFHRECTGMTESAYGLLTTEASAVWACDLCLKTKEIQSVYIREGMGQL
VAANDG

Figure 4



ClustalW Formatted Alignments






**Figure 6**

Figure 7





Figure 8

SEQUENCE LISTING PART

<110> The Genetics Company

<120> A new essential downstream component of the wingless signalling pathway

<130> P133559

<160> 10

<170> PatentIn version 3.1

<210> 1

<211> 2448

<212> DNA

<213> Drosophila melanogaster

<400> 1

```

atgacccaca atcttggtat ggcgccatat cgattgccgg gtccagcggg cggactctgt      60
ccgcccgaatt ttaagccgcc gcctcccacg gacatcatct cggcgccgag caatccgaag    120
aagcggcgaa aaacctcaag tgccgccaac tccgctgcag cgggtggctgc ggcggcggct    180
gcagcagctg ctgcgaattc catgcagcag cagcaggcgc caccacacc gcaggatttg    240
ctgccccctc cgccaatggg aggcttcgga gacaccatta ttgcctcgaa tccattcgac    300
gacagtcccc aggtgtcggc gatgtccagc tcagcggcgg cggcgatggc ggccatgaat    360
cagatgggcg gcggaccagg aggtggtcac tttggcggcg gtggaccggg tgggcacccg    420
cactgggaag accgcatggg catgggcggt ggacctctc cccgcctca catgcatccc    480
catatgcacc cgcacatccc aggcggacct atgggtcacc cacatggccc acatccgcac    540
atgggtggtc cacctccaat gcgaggaatg agcccatgc accccatca aatgggaccg    600
ggaccaggcg tcggactacc gccgcatatg aatcacggaa ggccaggggg acctggtggt    660
cctggaggac ccgtcccaat gggtagtccc atgggtggaa tagctggcat gggcggcatg    720
agcccaatgg gcggaatggg aggcgccagc atatcaccac atcacatggg catgggtggt    780
ctgtcgccca tgggaggcgg tcccaacgga cccaatccgc gagccatgca gggttcacccg    840
atgggcgggtc cggggcagaa ctgcgcaatg aactcactgc ctatgggttc gccaatgggc    900
aatccaattg gcagccggtt gggccctccc tcgggaccgg gccctgggaa tcccggcaat    960
accggcggac cacagcagca acaacaaca cctccgcagc caccgatgaa caacgggcag   1020
atgggtcctc ctctctgca cagtccgctc ggaaacggac caacgggtca tggcagtcac   1080
atgcctggag gaccaatccc aggaccaggt cctgggcctg gcggcctagt aggtcccggg   1140
ggcatctccc ccgcgcacgg caataacccg ggtggttctg ggaacaacat gctcggcggg   1200
aatcccggcg gcggcaacag caacaacaac ggaagcaata caagtaacgc cagcaacaac   1260
aatcaaaatc ctcacctctc gccagcagcc ggacgcctgg gagtgccgac gtcgatgcag   1320
tcgaatggac cttcggtatc atcggtagcc tcctcatcgg ttccctcgcc cgccacgccc   1380

```

acgctcacgc ccacatcgac ggccacgtcc atgtccacgt cagtgcctac atcctcgcca 1440
 gcgcgcgccg ccatgtcacc gcatcactcg ctaaacagcg ccggcccgag tccgggcatg 1500
 cccaactcgg gaccagccc gctgcagtca ccagccggac ccaatggccc caataacaac 1560
 aacagcaata acaacaacgg acccatgatg ggccagatga tcccgaacgc agttcctatg 1620
 cagcaccagc agcacatggg cggcggccca cctggccacg ggcccggacc aatgcccgga 1680
 atgggcatga accaaatgct gccaccgcag caaccctccc atcttggctc ccgcatccg 1740
 aatatgatga accacccgca tcacccgcac catcactctg gcggaccacc gccgcacatg 1800
 atgggtggac ccggaatgca cggcggctct gctggaatgc ctctcatat gggcggagga 1860
 cctaatacgc acatgatggg cggtcgcac gggaacgcgg gtccgcacat gggccacggc 1920
 cacatgggtg gagtaccagg tccaggaccc ggaccggcg gcatgaacgg acccccgcac 1980
 ccgcacatgt ccccgacca cggacatccg catcaccacc acaatccgat gggcggccca 2040
 ggtccaaata tggtcggcg tggtggagga ggtcccatgg gtcccgtgg accgatgggc 2100
 aacatggggc ccatgggagg tggcccgatg ggccggccca tggggtagg tcccagccg 2160
 atgacaatgg gcggcgggaa gatgtacccg ccgggacagc caatggctct taatccgcag 2220
 aaccggaatg cgcgcgccat atatccttgt ggcatgtgcc acaaggaggc gaacgacaac 2280
 gacgaagccg tggtctgtga atccggttgt aactttttct ttcacagaac ctgtgttggc 2340
 ctgacagagg cggccttcca aatgctcaac aaggagggtg ttgccgagtg gtgctgcgac 2400
 aagtgcgtgt cttccaagca tattcccatg gtcaagttca agtgttga 2448

<210> 2
 <211> 1366
 <212> DNA
 <213> Human

<400> 2
 ggatcccac atgcccgcg agaactctcc agtcccgcg tacaaagttt cctcgcatgg 60
 tgggtgatagt ggactggatg ggtaggagg accaggtgta caactaggaa gccagataa 120
 gaaaaagcgc aaggcaaata cacagggacc ttctttccct ccattgtctg agtatgctcc 180
 accaccgaat ccaaactctg accatctagt ggctgctaata ccatttgatg acaactataa 240
 tactatttcc tataaaccac taccttcgtc aaatccatat cttggccctg gttatcctgg 300
 ctttgagggc tatagtacat tcagaatgcc acctcacgtt ccccaagaa tgtcttcccc 360
 atactgtggg octtactcac tcaggaacca gccacacca tttctcaga atcctctggg 420
 catgggtttt aatcgacctc atgcttttaa ctttgggcca catgataatt caagtttcgg 480
 taatccatct tataataatg cactaagtca gaatgtcaac atgctaata aacattttag 540
 acaaaatcct gctgaaaatt tcagtcaaata tctccacag aatgctagcc aagtttctaa 600

ccccgatttg gcatctaatt ttgttcctgg aaataattca aattttactt ctccgtaga 660
 atctaatacat tcttttatct ctccccaaa cacttttggg caagcaaaag caccaccccc 720
 aaaacaagac ttactcaag gagcaaccaa aaacactaat caaaattcct ctgctcatcc 780
 acctcacttg aatatggatg acacagtga tcaagagtaatt attgaattaa aaaatgttaa 840
 tcgaaacaat gcagtaaatac aggagaacag ccgttcaagt agcactgaag ccacaaacaa 900
 taaccctgca aatgggacgc agaataagcc acgacaacca agagggtgag cagatgcctg 960
 caccacagaa aaaagcaata aatcctctct tcacccaaac cgtcatggcc attcgtcttc 1020
 tgaccagtg tctccttggt gaatttgtac aaacgagggt aacgatgatc aggatgccat 1080
 cttatgtgag gcctcttgct agaaatgggt tcatcggatc tgtactggaa tgactgaaac 1140
 agcttatggc ctcttaactg cagaagcatc tgcagtatgg ggctgtgata cctgtatggc 1200
 tgacaaagat gtccagttaa tgcgtactag agaaactttt ggtccatctg cagtgggcag 1260
 tgatgcttaa tcaaaggcat taactaaagt gggtttattt tcctgtgcat tgcagaagtt 1320
 cattgacaca ggattttaat gttttacatt attttttttaa atgcat 1366

<210> 3
 <211> 1449
 <212> DNA
 <213> Human

<400> 3
 cccgggtccc ccaactccatg gccgcctcgg cgcgcctccc accggacaag ctggagggag 60
 gtggcgggcc cgcaccgccc cctgcgcgcg ccagcaccgg gaggaagcag ggcaaggccg 120
 gtctgcaaat gaagagtcca gaaaagaagc gaagggaagtc aaataactcag ggccctgcat 180
 actcacatct gacggagttt gcaccacccc caactcccat ggtggatcac ctggttgcat 240
 ccaacccttt tgaagatgac ttcggagccc ccaaagtggg ggttgacagc cctccattcc 300
 ttggcagtc tgtgcccttc ggaggcttcc gtgtgcaggg gggcatggcg ggccaggtag 360
 ccccaggcta cagcactgga ggtggagggg gcccccagcc actccgtcga cagccacccc 420
 ccttccctcc caatcctatg ggccctgctt tcaacatgcc ccccagggt cctggctacc 480
 ccccccagg caacatgaac tttcccagcc aacccttcaa ccagcctctg ggtcaaaact 540
 ttagtccctc cagtgggcag atgatgccgg gccagtgagg gggatttggg cccatgatct 600
 caccacccat gggacagcct ccagagcag agctgggccc accttctctg tcccaacgat 660
 ttgctcagcc aggggctcct ttggccctt ctctctcca gagacctggg caggggctcc 720
 ccagcctgcc gcctaacaca agtccctttc ctgggtccgga cctgggcttt cctggccctg 780
 gtgggtgagga tggggggaag cccttgaatc cacctgcttc tactgctttt ccccaggagc 840
 cccactcagg ctccccggct gctgctgtta atgggaacca gccagtttc cccccgaaca 900

gcagtgggcg ggggtggggc actccagatg ccaacagctt ggcacccctt ggcaaggcag 960
 gtggggggctc cgggccccag cctccccag gcttggtgta cccatgtggt gcctgtcgga 1020
 gtgagggtgaa cgatgaccag gatgccattc tgtgtgaggc ctctgccag aaatgggtcc 1080
 accgtgagtg cacaggcatg actgagagcg cctatgggct gctgaccact gaagcttctg 1140
 ccgtctgggc ctgcatctc tgcctcaaga ccaaggagat ccagtctgtc tacatccgtg 1200
 agggcatggg gcagctgggtg gctgctaacg atgggtgacg ctgggtgaagt ggcccaggga 1260
 agtgcacatg tctctccctg ctctccaggt gtgatttttt tgatgtttgg ctcttggtcc 1320
 ttgtttccac tggctttcca tccccatggg gcagaaacag tggctcctgg gagcagaaaa 1380
 ggaattgagg tgggcaggca gaagagcctg gattgtcac tgttttggga aacttacatg 1440
 ttgagatct 1449

<210> 4
 <211> 1254
 <212> DNA
 <213> Mouse

<400> 4
 atgtcagcgg aacaggacaa ggagcccatc gcgctgaaga gagttagagg tgggtgacagt 60
 ggactggatg ggtaggagg gcccaatata caactcggaa gccagataa gaaaaaacgc 120
 aaggccaaca cacagggatc ttcctttcct ccgctgtcag agtacgcgcc accaccgaat 180
 ccaaactccg accatctagt ggctgccaat ccgttcgatg acagctacaa caccatttcc 240
 tataaaccac tgccttcac taatccatat cttggccctg ggtatcctgg ctttgagggc 300
 tacagcacat tcagaatgcc accccacgtc cctccaagaa tgtcttctcc ctactgtggt 360
 ccttactcac tcaggaatca gcccaccca tttcctcaga atccgttggg catgggcttt 420
 aaccggcctc atgcttttaa ctttgggccg catgataatt cgaattttgg aaaccacct 480
 tataataatg tactgactca ggacattaac atgcccggtc agcatttttag acaaggctct 540
 gctgaaaact tcagtcagat tccccgcag aatgttggtc aagtgtctaa ccctgacctc 600
 gcatctaatt ttgcccttg gaataattca aattttacct ctccgttaga aacgaatcat 660
 tcgtttatcc ccccccaaa cgcgtttggc caagcaaaag ctccacttcc caaacaagac 720
 ttcactcaag gggcaacaa aacccgaat cagaattcgt ccactcacc acctcaccta 780
 aatatggagg atccagtcaa tcagagtaac gtcaggttaa aaaatgtcaa cagaaacaac 840
 gttgtccaag agaacagccg ttcgggcagc gcagaggcca ccaacaacca tgcgaatggg 900
 acccagaaca agccccggca gccaggggc gcagctgacc tgtgcacccc cgacaaaagc 960
 cgcaagttct ccctgtctcc cagccggcat ggccattcct cctctgacct tgtgtacctg 1020
 tgcgggattt gtacaaatga agtgaatgac gatcaggacg ccattctgtg tgaagcctct 1080

tgtcagaagt ggtttcatcg catctgcact ggaatgaccg aaacagccta cgggctcctg 1140
 acagcggaag catccgcagt gtggggctgt gacacgtgca tggctgacaa ggatgtccag 1200
 ctcatgcgca ctagagagggc ctttggtcca cctgccgtgg gcggcgatgc ctaa 1254

<210> 5
 <211> 1221
 <212> DNA
 <213> Mouse

<400> 5
 atggccgcct cggcgccgcc cccaccggac aagctggagg gaggcagcgg ccccgaccg 60
 cccccgcgc cgcacagcac cgggaggaag cagggcaagg ccggtctgca aatgaagagc 120
 cctgaaaaga agcgaagaaa gtccaatact cagggtcctg catattcaca tctgacggag 180
 ttgcgccac ccccgacccc catggtggat catctggtcg cttctaacc ttttgaggat 240
 gatttcggag ccctaaagt ggggggcgca ggccctcgt tctcggcag tccggtgccc 300
 tttggaggct ttcgtgtaca ggggggcatg gcaggccagg taccaccaag ctacggcact 360
 ggaggaggag ggggtcccca gccacttcgt cggcagcccc ctcttttcc cccagccct 420
 atgggtccag cttttaatat gcccctcag ggtccctggg gtaccccgcc cctggcaac 480
 atgaactttc ccagtcaacc cttcaaccag tctctgggcc aaaactttag ccacctggt 540
 gggcaggtga tgccaggccc agtaggcgga tttggtccca tgatctcacc gaccatggga 600
 cagcctccta gaggggagct gggctctcct cctctcccc aacgctttac ccaaccagga 660
 gcaccttatg gtctttctct tcagagacct ggtcagggac tcaccagct gccctccaac 720
 acaagtccct tccctggtcc agaccctggt tttcctggac ctggcggtga ggatggtggg 780
 aagcccttga acccaccggc tcccaccgcc tttccccagg aagcaccatt cgggctcccc 840
 gctgctgctg tcaatgggaa tcagcccagt tcccccccta gcagcagtg tggaggtggg 900
 ggcactccag atgccaacag tctggcacc cccggcaagg cagggggagg ctgagggccc 960
 cagcctcccc caggcctggt gtaccctgc ggtgcctgcc gtagtgaggt aaatgatgac 1020
 caggatgcca ttctgtgtga ggctcctgc cagaagtggg ttcaccgga gtgcaccggc 1080
 atgaccgaga gtgcctacgg cctgctgacc accgaggcct ctgccgtctg ggctgtgat 1140
 ctttgccca agaccaagga gatccagtct gtctacatcc gagagggcat gggccagttg 1200
 gtggctgcta acgatgggtg a 1221

<210> 6
 <211> 815
 <212> PRT
 <213> *Drosophila melanogaster*

<400> 6

Met Thr His Asn Leu Gly Met Ala Pro Tyr Arg Leu Pro Gly Pro Ala
1 5 10 15

Gly Gly Leu Cys Pro Pro Asp Phe Lys Pro Pro Pro Pro Thr Asp Ile
20 25 30

Ile Ser Ala Pro Ser Asn Pro Lys Lys Arg Arg Lys Thr Ser Ser Ala
35 40 45

Ala Asn Ser Ala Ala Ala Val Ala Ala Ala Ala Ala Ala Ala Ala Ala
50 55 60

Ala Asn Ser Met Gln Gln Gln Gln Ala Pro Pro Thr Pro Gln Asp Leu
65 70 75 80

Leu Pro Pro Pro Pro Met Gly Gly Phe Gly Asp Thr Ile Ile Ala Ser
85 90 95

Asn Pro Phe Asp Asp Ser Pro Gln Val Ser Ala Met Ser Ser Ser Ala
100 105 110

Ala Ala Ala Met Ala Ala Met Asn Gln Met Gly Gly Gly Pro Gly Gly
115 120 125

Gly His Phe Gly Gly Gly Gly Pro Gly Gly His Pro His Trp Glu Asp
130 135 140

Arg Met Gly Met Gly Gly Gly Pro Pro Pro Pro Pro His Met His Pro
145 150 155 160

His Met His Pro His His Pro Gly Gly Pro Met Gly His Pro His Gly
165 170 175

Pro His Pro His Met Gly Gly Pro Pro Pro Met Arg Gly Met Ser Pro
180 185 190

Met His Pro His Gln Met Gly Pro Gly Pro Gly Val Gly Leu Pro Pro
195 200 205

His Met Asn His Gly Arg Pro Gly Gly Pro Gly Gly Pro Gly Gly Pro
210 215 220

Val Pro Met Gly Ser Pro Met Gly Gly Ile Ala Gly Met Gly Gly Met
225 230 235 240

Ser Pro Met Gly Gly Met Gly Gly Pro Ser Ile Ser Pro His His Met
245 250 255

Gly Met Gly Gly Leu Ser Pro Met Gly Gly Gly Pro Asn Gly Pro Asn
 260 265 270

Pro Arg Ala Met Gln Gly Ser Pro Met Gly Gly Pro Gly Gln Asn Ser
 275 280 285

Pro Met Asn Ser Leu Pro Met Gly Ser Pro Met Gly Asn Pro Ile Gly
 290 295 300

Ser Pro Leu Gly Pro Pro Ser Gly Pro Gly Pro Gly Asn Pro Gly Asn
 305 310 315 320

Thr Gly Gly Pro Gln Gln Gln Gln Gln Gln Pro Pro Gln Pro Pro Met
 325 330 335

Asn Asn Gly Gln Met Gly Pro Pro Pro Leu His Ser Pro Leu Gly Asn
 340 345 350

Gly Pro Thr Gly His Gly Ser His Met Pro Gly Gly Pro Ile Pro Gly
 355 360 365

Pro Gly Pro Gly Pro Gly Gly Leu Val Gly Pro Gly Gly Ile Ser Pro
 370 375 380

Ala His Gly Asn Asn Pro Gly Gly Ser Gly Asn Asn Met Leu Gly Gly
 385 390 395 400

Asn Pro Gly Gly Gly Asn Ser Asn Asn Asn Gly Ser Asn Thr Ser Asn
 405 410 415

Ala Ser Asn Asn Asn Gln Asn Pro His Leu Ser Pro Ala Ala Gly Arg
 420 425 430

Leu Gly Val Pro Thr Ser Met Gln Ser Asn Gly Pro Ser Val Ser Ser
 435 440 445

Val Ala Ser Ser Ser Val Pro Ser Pro Ala Thr Pro Thr Leu Thr Pro
 450 455 460

Thr Ser Thr Ala Thr Ser Met Ser Thr Ser Val Pro Thr Ser Ser Pro
 465 470 475 480

Ala Pro Pro Ala Met Ser Pro His His Ser Leu Asn Ser Ala Gly Pro
 485 490 495

Ser Pro Gly Met Pro Asn Ser Gly Pro Ser Pro Leu Gln Ser Pro Ala

500					505					510					
Gly	Pro	Asn	Gly	Pro	Asn	Asn	Asn	Asn	Ser	Asn	Asn	Asn	Asn	Gly	Pro
		515					520					525			
Met	Met	Gly	Gln	Met	Ile	Pro	Asn	Ala	Val	Pro	Met	Gln	His	Gln	Gln
		530					535					540			
His	Met	Gly	Gly	Gly	Pro	Pro	Gly	His	Gly	Pro	Gly	Pro	Met	Pro	Gly
		545					550					555			560
Met	Gly	Met	Asn	Gln	Met	Leu	Pro	Pro	Gln	Gln	Pro	Ser	His	Leu	Gly
				565					570					575	
Pro	Pro	His	Pro	Asn	Met	Met	Asn	His	Pro	His	His	Pro	His	His	His
				580					585					590	
Pro	Gly	Gly	Pro	Pro	Pro	His	Met	Met	Gly	Gly	Pro	Gly	Met	His	Gly
				595					600					605	
Gly	Pro	Ala	Gly	Met	Pro	Pro	His	Met	Gly	Gly	Gly	Pro	Asn	Pro	His
				610					615					620	
Met	Met	Gly	Gly	Pro	His	Gly	Asn	Ala	Gly	Pro	His	Met	Gly	His	Gly
				625					630					635	640
His	Met	Gly	Gly	Val	Pro	Gly	Pro	Gly	Pro	Gly	Pro	Gly	Gly	Met	Asn
				645					650					655	
Gly	Pro	Pro	His	Pro	His	Met	Ser	Pro	His	His	Gly	His	Pro	His	His
				660					665					670	
His	His	Asn	Pro	Met	Gly	Gly	Pro	Gly	Pro	Asn	Met	Phe	Gly	Gly	Gly
				675					680					685	
Gly	Gly	Gly	Pro	Met	Gly	Pro	Gly	Gly	Pro	Met	Gly	Asn	Met	Gly	Pro
				690					695					700	
Met	Gly	Gly	Gly	Pro	Met	Gly	Gly	Pro	Met	Gly	Val	Gly	Pro	Lys	Pro
				705					710					715	720
Met	Thr	Met	Gly	Gly	Gly	Lys	Met	Tyr	Pro	Pro	Gly	Gln	Pro	Met	Val
				725					730					735	
Phe	Asn	Pro	Gln	Asn	Pro	Asn	Ala	Pro	Pro	Ile	Tyr	Pro	Cys	Gly	Met
				740					745					750	

Cys His Lys Glu Val Asn Asp Asn Asp Glu Ala Val Phe Cys Glu Ser
755 760 765

Gly Cys Asn Phe Phe Phe His Arg Thr Cys Val Gly Leu Thr Glu Ala
770 775 780

Ala Phe Gln Met Leu Asn Lys Glu Val Phe Ala Glu Trp Cys Cys Asp
785 790 795 800

Lys Cys Val Ser Ser Lys His Ile Pro Met Val Lys Phe Lys Cys
805 810 815

<210> 7
<211> 419
<212> PRT
<213> Human

<400> 7

Met Pro Ala Glu Asn Ser Pro Ala Pro Ala Tyr Lys Val Ser Ser His
1 5 10 15

Gly Gly Asp Ser Gly Leu Asp Gly Leu Gly Gly Pro Gly Val Gln Leu
20 25 30

Gly Ser Pro Asp Lys Lys Lys Arg Lys Ala Asn Thr Gln Gly Pro Ser
35 40 45

Phe Pro Pro Leu Ser Glu Tyr Ala Pro Pro Pro Asn Pro Asn Ser Asp
50 55 60

His Leu Val Ala Ala Asn Pro Phe Asp Asp Asn Tyr Asn Thr Ile Ser
65 70 75 80

Tyr Lys Pro Leu Pro Ser Ser Asn Pro Tyr Leu Gly Pro Gly Tyr Pro
85 90 95

Gly Phe Gly Gly Tyr Ser Thr Phe Arg Met Pro Pro His Val Pro Pro
100 105 110

Arg Met Ser Ser Pro Tyr Cys Gly Pro Tyr Ser Leu Arg Asn Gln Pro
115 120 125

His Pro Phe Pro Gln Asn Pro Leu Gly Met Gly Phe Asn Arg Pro His
130 135 140

Ala Phe Asn Phe Gly Pro His Asp Asn Ser Ser Phe Gly Asn Pro Ser
145 150 155 160

10

Tyr Asn Asn Ala Leu Ser Gln Asn Val Asn Met Pro Asn Gln His Phe
 165 170 175

Arg Gln Asn Pro Ala Glu Asn Phe Ser Gln Ile Pro Pro Gln Asn Ala
 180 185 190

Ser Gln Val Ser Asn Pro Asp Leu Ala Ser Asn Phe Val Pro Gly Asn
 195 200 205

Asn Ser Asn Phe Thr Ser Pro Leu Glu Ser Asn His Ser Phe Ile Pro
 210 215 220

Pro Pro Asn Thr Phe Gly Gln Ala Lys Ala Pro Pro Pro Lys Gln Asp
 225 230 235 240

Phe Thr Gln Gly Ala Thr Lys Asn Thr Asn Gln Asn Ser Ser Ala His
 245 250 255

Pro Pro His Leu Asn Met Asp Asp Thr Val Asn Gln Ser Asn Ile Glu
 260 265 270

Leu Lys Asn Val Asn Arg Asn Asn Ala Val Asn Gln Glu Asn Ser Arg
 275 280 285

Ser Ser Ser Thr Glu Ala Thr Asn Asn Asn Pro Ala Asn Gly Thr Gln
 290 295 300

Asn Lys Pro Arg Gln Pro Arg Gly Ala Ala Asp Ala Cys Thr Thr Glu
 305 310 315 320

Lys Ser Asn Lys Ser Ser Leu His Pro Asn Arg His Gly His Ser Ser
 325 330 335

Ser Asp Pro Val Tyr Pro Cys Gly Ile Cys Thr Asn Glu Val Asn Asp
 340 345 350

Asp Gln Asp Ala Ile Leu Cys Glu Ala Ser Cys Gln Lys Trp Phe His
 355 360 365

Arg Ile Cys Thr Gly Met Thr Glu Thr Ala Tyr Gly Leu Leu Thr Ala
 370 375 380

Glu Ala Ser Ala Val Trp Gly Cys Asp Thr Cys Met Ala Asp Lys Asp
 385 390 395 400

Val Gln Leu Met Arg Thr Arg Glu Thr Phe Gly Pro Ser Ala Val Gly
 405 410 415

Ser Asp Ala

<210> 8
 <211> 406
 <212> PRT
 <213> Human

<400> 8

Met Ala Ala Ser Ala Pro Pro Pro Pro Asp Lys Leu Glu Gly Gly Gly
 1 5 10 15

Gly Pro Ala Pro Pro Pro Ala Pro Pro Ser Thr Gly Arg Lys Gln Gly
 20 25 30

Lys Ala Gly Leu Gln Met Lys Ser Pro Glu Lys Lys Arg Arg Lys Ser
 35 40 45

Asn Thr Gln Gly Pro Ala Tyr Ser His Leu Thr Glu Phe Ala Pro Pro
 50 55 60

Pro Thr Pro Met Val Asp His Leu Val Ala Ser Asn Pro Phe Glu Asp
 65 70 75 80

Asp Phe Gly Ala Pro Lys Val Gly Val Ala Ala Pro Pro Phe Leu Gly
 85 90 95

Ser Pro Val Pro Phe Gly Gly Phe Arg Val Gln Gly Gly Met Ala Gly
 100 105 110

Gln Val Pro Pro Gly Tyr Ser Thr Gly Gly Gly Gly Gly Pro Gln Pro
 115 120 125

Leu Arg Arg Gln Pro Pro Pro Phe Pro Pro Asn Pro Met Gly Pro Ala
 130 135 140

Phe Asn Met Pro Pro Gln Gly Pro Gly Tyr Pro Pro Pro Gly Asn Met
 145 150 155 160

Asn Phe Pro Ser Gln Pro Phe Asn Gln Pro Leu Gly Gln Asn Phe Ser
 165 170 175

Pro Pro Ser Gly Gln Met Met Pro Gly Pro Val Gly Gly Phe Gly Pro
 180 185 190

Met Ile Ser Pro Thr Met Gly Gln Pro Pro Arg Ala Glu Leu Gly Pro
 195 200 205

Pro Ser Leu Ser Gln Arg Phe Ala Gln Pro Gly Ala Pro Phe Gly Pro
 210 215 220

Ser Pro Leu Gln Arg Pro Gly Gln Gly Leu Pro Ser Leu Pro Pro Asn
 225 230 235 240

Thr Ser Pro Phe Pro Gly Pro Asp Pro Gly Phe Pro Gly Pro Gly Gly
 245 250 255

Glu Asp Gly Gly Lys Pro Leu Asn Pro Pro Ala Ser Thr Ala Phe Pro
 260 265 270

Gln Glu Pro His Ser Gly Ser Pro Ala Ala Ala Val Asn Gly Asn Gln
 275 280 285

Pro Ser Phe Pro Pro Asn Ser Ser Gly Arg Gly Gly Gly Thr Pro Asp
 290 295 300

Ala Asn Ser Leu Ala Pro Pro Gly Lys Ala Gly Gly Gly Ser Gly Pro
 305 310 315 320

Gln Pro Pro Pro Gly Leu Val Tyr Pro Cys Gly Ala Cys Arg Ser Glu
 325 330 335

Val Asn Asp Asp Gln Asp Ala Ile Leu Cys Glu Ala Ser Cys Gln Lys
 340 345 350

Trp Phe His Arg Glu Cys Thr Gly Met Thr Glu Ser Ala Tyr Gly Leu
 355 360 365

Leu Thr Thr Glu Ala Ser Ala Val Trp Ala Cys Asp Leu Cys Leu Lys
 370 375 380

Thr Lys Glu Ile Gln Ser Val Tyr Ile Arg Glu Gly Met Gly Gln Leu
 385 390 395 400

Val Ala Ala Asn Asp Gly
 405

<210> 9
 <211> 417
 <212> PRT
 <213> Mouse

<400> 9

Met Ser Ala Glu Gln Asp Lys Glu Pro Ile Ala Leu Lys Arg Val Arg
 1 5 10 15

Gly Gly Asp Ser Gly Leu Asp Gly Leu Gly Gly Pro Asn Ile Gln Leu
 20 25 30

Gly Ser Pro Asp Lys Lys Lys Arg Lys Ala Asn Thr Gln Gly Ser Ser
 35 40 45

Phe Pro Pro Leu Ser Glu Tyr Ala Pro Pro Pro Asn Pro Asn Ser Asp
 50 55 60

His Leu Val Ala Ala Asn Pro Phe Asp Asp Ser Tyr Asn Thr Ile Ser
 65 70 75 80

Tyr Lys Pro Leu Pro Ser Ser Asn Pro Tyr Leu Gly Pro Gly Tyr Pro
 85 90 95

Gly Phe Gly Gly Tyr Ser Thr Phe Arg Met Pro Pro His Val Pro Pro
 100 105 110

Arg Met Ser Ser Pro Tyr Cys Gly Pro Tyr Ser Leu Arg Asn Gln Pro
 115 120 125

His Pro Phe Pro Gln Asn Pro Leu Gly Met Gly Phe Asn Arg Pro His
 130 135 140

Ala Phe Asn Phe Gly Pro His Asp Asn Ser Asn Phe Gly Asn Pro Pro
 145 150 155 160

Tyr Asn Asn Val Leu Thr Gln Asp Ile Asn Met Pro Gly Gln His Phe
 165 170 175

Arg Gln Gly Ser Ala Glu Asn Phe Ser Gln Ile Pro Pro Gln Asn Val
 180 185 190

Gly Gln Val Ser Asn Pro Asp Leu Ala Ser Asn Phe Ala Pro Gly Asn
 195 200 205

Asn Ser Asn Phe Thr Ser Pro Leu Glu Thr Asn His Ser Phe Ile Pro
 210 215 220

Pro Pro Asn Ala Phe Gly Gln Ala Lys Ala Pro Leu Pro Lys Gln Asp
 225 230 235 240

Phe Thr Gln Gly Ala Thr Lys Thr Pro Asn Gln Asn Ser Ser Thr His
 245 250 255

Pro Pro His Leu Asn Met Glu Asp Pro Val Asn Gln Ser Asn Val Glu
 260 265 270

14

Leu Lys Asn Val Asn Arg Asn Asn Val Val Gln Glu Asn Ser Arg Ser
 275 280 285

Gly Ser Ala Glu Ala Thr Asn Asn His Ala Asn Gly Thr Gln Asn Lys
 290 295 300

Pro Arg Gln Pro Arg Gly Ala Ala Asp Leu Cys Thr Pro Asp Lys Ser
 305 310 315 320

Arg Lys Phe Ser Leu Leu Pro Ser Arg His Gly His Ser Ser Ser Asp
 325 330 335

Pro Val Tyr Pro Cys Gly Ile Cys Thr Asn Glu Val Asn Asp Asp Gln
 340 345 350

Asp Ala Ile Leu Cys Glu Ala Ser Cys Gln Lys Trp Phe His Arg Ile
 355 360 365

Cys Thr Gly Met Thr Glu Thr Ala Tyr Gly Leu Leu Thr Ala Glu Ala
 370 375 380

Ser Ala Val Trp Gly Cys Asp Thr Cys Met Ala Asp Lys Asp Val Gln
 385 390 395 400

Leu Met Arg Thr Arg Glu Ala Phe Gly Pro Pro Ala Val Gly Gly Asp
 405 410 415

Ala

<210> 10
 <211> 406
 <212> PRT
 <213> Mouse

<400> 10

Met Ala Ala Ser Ala Pro Pro Pro Pro Asp Lys Leu Glu Gly Gly Ser
 1 5 10 15

Gly Pro Ala Pro Pro Pro Ala Pro Pro Ser Thr Gly Arg Lys Gln Gly
 20 25 30

Lys Ala Gly Leu Gln Met Lys Ser Pro Glu Lys Lys Arg Arg Lys Ser
 35 40 45

Asn Thr Gln Gly Pro Ala Tyr Ser His Leu Thr Glu Phe Ala Pro Pro
 50 55 60

Pro Thr Pro Met Val Asp His Leu Val Ala Ser Asn Pro Phe Glu Asp
 65. 70 75 80

Asp Phe Gly Ala Pro Lys Val Gly Gly Ala Gly Pro Pro Phe Leu Gly
 85 90 95

Ser Pro Val Pro Phe Gly Gly Phe Arg Val Gln Gly Gly Met Ala Gly
 100 105 110

Gln Val Pro Pro Ser Tyr Gly Thr Gly Gly Gly Gly Gly Pro Gln Pro
 115 120 125

Leu Arg Arg Gln Pro Pro Pro Phe Pro Pro Ser Pro Met Gly Pro Ala
 130 135 140

Phe Asn Met Pro Pro Gln Gly Pro Trp Gly Thr Pro Pro Pro Gly Asn
 145 150 155 160

Met Asn Phe Pro Ser Gln Pro Phe Asn Gln Ser Leu Gly Gln Asn Phe
 165 170 175

Ser Pro Pro Gly Gly Gln Val Met Pro Gly Pro Val Gly Gly Phe Gly
 180 185 190

Pro Met Ile Ser Pro Thr Met Gly Gln Pro Pro Arg Gly Glu Leu Gly
 195 200 205

Pro Pro Pro Leu Pro Gln Arg Phe Thr Gln Pro Gly Ala Pro Tyr Gly
 210 215 220

Pro Ser Leu Gln Arg Pro Gly Gln Gly Leu Thr Gln Leu Pro Ser Asn
 225 230 235 240

Thr Ser Pro Phe Pro Gly Pro Asp Pro Gly Phe Pro Gly Pro Gly Gly
 245 250 255

Glu Asp Gly Gly Lys Pro Leu Asn Pro Pro Ala Pro Thr Ala Phe Pro
 260 265 270

Gln Glu Ala Pro Phe Gly Leu Pro Ala Ala Ala Val Asn Gly Asn Gln
 275 280 285

Pro Ser Phe Pro Pro Ser Ser Ser Gly Arg Gly Gly Gly Thr Pro Asp
 290 295 300

Ala Asn Ser Leu Ala Pro Pro Gly Lys Ala Gly Gly Gly Ser Gly Pro

[illegible]

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 October 2002 (03.10.2002)

PCT

(10) International Publication Number
WO 02/077023 A3

(51) International Patent Classification⁷: **C07K 14/435**,
C12N 15/12, C12Q 1/68, G01N 33/50, A61K 38/18,
C07K 16/18

(21) International Application Number: PCT/CH02/00063

(22) International Filing Date: 1 February 2002 (01.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/277,976 23 March 2001 (23.03.2001) US

(71) Applicant (for all designated States except US): **UNIVERSITÄT ZÜRICH** [CH/CH]; Rämistrasse 71, CH-8006 Zürich (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KRAMPS, Thomas** [DE/CH]; Dorfstrasse 51, CH-8037 Zürich (CH). **BASLER, Konrad** [CH/CH]; Traubenweg 34, CH-8700 Küsnacht (CH).

(74) Agent: **LIEBETANZ, Michael**; Isler & Pedrazzini AG, Postfach 6940, CH-8023 Zürich (CH).

(81) Designated States (national): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,

CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
16 January 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A NEW ESSENTIAL DOWNSTREAM COMPONENT OF THE WINGLESS SIGNALLING PATHWAY

(57) Abstract: The present invention relates to a new essential downstream component of the wingless signaling pathway. In particular, the invention relates to nucleotide sequences of the *Drosophila melanogaster* daughter of legless (*doll*) gene, of its encoded proteins, as well as derivatives, fragments and analogues thereof. The invention includes vertebrate and invertebrate homologues of the *Doll* protein, comprising proteins that contain a stretch of amino acids with similarity to the *Drosophila Doll* gene. Methods for producing the *Doll* protein, derivatives and analogs, e.g. by recombinant means, and antibodies to *Doll* are provided by the present invention as well. The invention also relates to methods for performing high throughput screening assays for compounds modulating *Doll* function in the Wnt pathway.

INTERNATIONAL SEARCH REPORT

Intern. application No	PCT/CH 02/00063
------------------------	-----------------

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 C07K14/435 C12N15/12 C12Q1/68 G01N33/50 A61K38/18
 C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, EMBL, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! EBI; 18 August 1999 (1999-08-18) CELNIKER ET AL.: "Drosophila melanogaster chromosome 3R, region 100c, clone BACR10009" Database accession no. AC009349 XP002217662 100% identity with SEQ ID NO:1 in 2327 of 2448 nt (2327-1: 140498-142824) the whole document <div style="text-align: center; margin-top: 20px;">-/--</div>	1-4, 28-32

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

22 October 2002

Date of mailing of the international search report

06/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Niebuhr-Ebel, K

INTERNATIONAL SEARCH REPORT

Intern: Application No

PCT/CH 02/00063

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! EBI; 4 November 1999 (1999-11-04) MUZNY ET AL.: "Homo sapiens chromosome 3 clone RP1-458H3" Database accession no. AC012674 XP002217663 99.8% (100% ungapped) identity with SEQ ID NO:2 in 1226 of 1366 nt (143-1366: 118972-120197) the whole document</p>	1,2,5,6, 28-32
X	<p>DATABASE EMBL 'Online! EBI; 24 January 2000 (2000-01-24) MUZNY ET AL.: "Homo sapiens chromosome 3 clone RP11-498A2" Database accession no. AC021890 XP002217664 99.9% (100% ungapped) identity with SEQ ID NO:3 in 1285 of 1449 nt (1449-165: 130620-131903) 87.1% (88% ungapped) identity with SEQ ID No:5 in 1079 of 1221 nt (1221-148: 130831-131903) the whole document</p>	1,2,7,8, 11,12, 28-32
X	<p>DATABASE EMBL 'Online! EBI; 8 February 2001 (2001-02-08) ADACHI ET AL.: "Mus musculus 10 days embryo whole body cDNA, clone 2600014C22: PHD-finger containing protein" Database accession no. AK011208 XP002217665 100% identity with SEQ ID NO:4 in 1254 of 1254 nt (1-1254: 265-1518) the whole document</p>	1,2,9, 10,28-32
P,X	<p>DATABASE EMBL 'Online! EBI; 11 August 2001 (2001-08-11) NI ET AL.: "Mus musculus chromosome 3 clone rp23-418c12 strain C57BL/6J" Database accession no. AC104329 XP002217666 98.7% (99.3% ungapped) with SEQ ID NO:5 in 1073 of 1221 nt (151-1221:86821-87888) the whole document</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1,2,11, 12,28-32

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/CH 02/00063

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	DATABASE EMBL 'Online! EBI; 11 October 2001 (2001-10-11) DRMANAC ET AL.: "Novel human diagnostic protein #27658" Database accession no. ABG27667 XP002217667 95.3% (95.3% ungapped) identity with SEQ ID No:7 over entire length the whole document & WO 01 75067 A (HYSEQ INC.) 11 October 2001 (2001-10-11) SEQ ID No: 58026 claim 20	13,14, 17,18, 38,48-50
A	MOON R T ET AL: "WNTs modulate cell fate and behavior during vertebrate development" TRENDS IN GENETICS, ELSEVIER SCIENCE PUBLISHERS B.V. AMSTERDAM, NL, vol. 13, no. 4, 1 April 1997 (1997-04-01), pages 157-162, XP004056903 ISSN: 0168-9525 the whole document	

INTERNATIONAL SEARCH REPORT

Int'l application No.
PCT/CH 02/00063

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1, 2, 13, 14, 33, 34, 36, 38-48, 50-54 (partially), 25 (completely)

Present claims 1, 2, 13, 14, 33, 34, 36, 38-48 and 50-54 relate to "Daughter of Legless" (DOLL) genes or proteins or to "DHD" domains without technically specifying said genes or proteins by referring to a SEQ ID. DOLL and "DHD domain" are arbitrary denominations and do not have a meaning for a person skilled in the art. Consequently such arbitrary names are not considered a technical feature. A gene or protein (domain) represents a chemical compound and should therefore be defined by structural features such as a SEQ ID or a certain degree of homology thereto. Consequently the search with respect to claims 1, 2, 13, 14, 33, 34, 36, 38-48 and 50-54 has been restricted to SEQ ID NOs: 1-10.

Claim 25 relates to a nucleic acid sequence coding for a peptide comprising a "low overall amino acid sequence identity" and "conserved domains" with a sequence identity of 50-100%. Without a clearer definition of said conserved domains in terms of sequence, length or position within the protein, no meaningful search is possible for said claim.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: application No

PCT/CH 02/00063

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0175067	A	11-10-2001	AU 3486501 A	14-08-2001
			AU 3995501 A	12-09-2001
			AU 4925101 A	15-10-2001
			AU 4972801 A	15-10-2001
			AU 5119401 A	15-10-2001
			AU 5521401 A	15-10-2001
			WO 0157266 A1	09-08-2001
			WO 0164839 A2	07-09-2001
			WO 0175067 A2	11-10-2001
			WO 0175064 A2	11-10-2001
			WO 0174836 A1	11-10-2001
			WO 0175093 A1	11-10-2001
			US 6436703 B1	20-08-2002
			US 2002061567 A1	23-05-2002
			US 2002146757 A1	10-10-2002

Form PCT/ISA/210 (patent family annex) (July 1992)

THIS PAGE BLANK (USPTO)